

JCO41 Rec'd PCT/PTO 31 MAY 2001
09/857067

PCT
\$



TRANSMITTAL LETTER TO THE UNITED STATES

ATTORNEY'S DOCKET NUMBER 50555

DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371

U.S. APPLICATION NO. (If known, see 37 CFR 1.5)

INTERNATIONAL APPLICATION NO.	INTERNATIONAL FILING DATE	PRIORITY DATE CLAIMED
PCT/EP 99/09413	2 December 1999	6 August 1999 4 December 1998

TITLE OF INVENTION: 3-(HETEROCYCLYL)-SUBSTITUTED BENZOYLPYRAZOLES

APPLICANT(S) FOR DO/EO/US Ulf NEIDLEIN, Norbert GOETZ, Ernst BAUMANN, Wolfgang von DEYN, Steffen KUDIS,
Klaus LANGEMANN, Guido MAYER, Ulf MISSLITZ, Matthias WITSCHKEI,
Martina OTTEN, Karl-Otto WESTPHALEN, Helmut WALTER

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a FIRST submission of items concerning a filing under 35 U.S.C. 371.
 2. ☐ This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371.
 3. ☒ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
 4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
 5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2)).
 - a. ☒ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ has been transmitted by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/USO).
 6. ☒ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
 7. ☐ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)).
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☐ have not been made and will not be made.
 8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
 9. ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
 10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).
- Items 11. to 16. below concern other document(s) or information included:
11. ☒ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
 12. ☒ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
 13. ☒ A FIRST preliminary amendment.
☐ A SECOND or SUBSEQUENT preliminary amendment.
 14. ☐ A substitute specification.
 15. ☐ A change of power of attorney and/or address letter.
 16. ☒ Other items or information.
International Search Report
International Preliminary Examination Report

U.S. Appl. No. (If Known) INTERNATIONAL APPLN. NO.
PCT/EP99/09413ATTORNEY'S DOCKET NO.
50555

	CALCULATIONS	PTO USE ONLY
17. /X/ The following fees are submitted		
BASIC NATIONAL FEE (37 CFR 1.492(a)(1)-(5)):		
Search Report has been prepared by the		
EPO or JPO.....\$860.00	860.00	
International preliminary examination fee paid to USPTO		
(37 CFR 1.482).....\$750.00		
No international preliminary examination fee paid to		
USPTO (37 CFR 1.482) but international search fee paid		
to USPTO (37 CFR 1.445(a)(2)).....\$700.00		
Neither international preliminary examination fee		
(37 CFR 1.482) nor international search fee		
(37 CFR 1.445(a)(2)) paid to USPTO\$ 970.00		
International preliminary examination fee paid to		
USPTO (37 CFR 1.482) and all claims satisfied pro		
-visions of PCT Article 33(2)-(4).....\$96.00		
ENTER APPROPRIATE BASIC FEE AMOUNT = \$ 860.00		
Surcharge of \$130.00 for furnishing the oath or declaration		
later than // 20 // 30 months from the earliest		
claimed priority date (37 CFR 1.492(e)).		
Claims	Number Filed	Number Extra
Total Claims	11 -20	
Indep. Claims	1 -3	
Multiple dependent claim(s) (if applicable)		+270.
TOTAL OF ABOVE CALCULATION		= 860.00
Reduction of 1/2 for filing by small entity, if applicable.		
Verified Small Entity statement must also be filed		
(Note 37 CFR 1.9, 1.27, 1.28).		
SUBTOTAL		= 860.00
Processing fee of \$130. for furnishing the English		
translation later than // 20 // 30 months from the		
earliest claimed priority date (37 CFR 1.492(f)).		
TOTAL NATIONAL FEE		= 860.00
Fee for recording the enclosed assignment (37 CFR 1.21(h)).		
The assignment must be accompanied by an appropriate cover		
sheet (37 CFR 3.28, 3.31) \$40.00 per property		
TOTAL FEES ENCLOSED		= \$ 900.00
Amount to be		
refunded: \$		
Charged \$		

a./X/ A check in the amount of \$ 900. to cover the above fees is enclosed.

b./ / Please charge my Deposit Account No. _____ in the amount of \$ _____ to cover the above fees. A duplicate copy of this sheet is enclosed.

c./X/ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 11-0345. A duplicate copy of this sheet is enclosed.**NOTE:** Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b) must be filed and granted to restore the application to pending status.**SEND ALL CORRESPONDENCE TO:**KEIL & WEINKAUF
1101 Connecticut Ave., N.W.
Washington, D. C. 20036

SIGNATURE

Herbert B. Keil

NAME

Registration No. 18,967

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re the Application of)
NEIDLEIN et al.) BOX PCT

International Application)
PCT/EP 99/09413)

Filed: December 2, 1999)

For: 3-(HETEROCYCLYL)-SUBSTITUTED BENZOYLPYRAZOLES

PRELIMINARY AMENDMENT

Honorable Commissioner of
Patents and Trademarks
Washington, D.C. 20231

Sir:

Prior to examination, kindly amend the above-identified application as follows:

IN THE CLAIMS

Cancel claims 11 and 13.

Please amend the claims as shown in the attached sheet.

REMARKS

The claims have been amended to eliminate multiple dependency and to put them in better form for U.S. filing. No new matter is included. A clean copy of the claims is attached.

Favorable action is solicited.

Respectfully submitted,

KEIL & WEINKAUF



Herbert B. Keil
Reg. No. 18,967

1101 Connecticut Ave., N.W.
Washington, D.C. 20036

(202)659-0100

PRELIMINARY AMENDMENTS TO THE CLAIMS - OZ 50555

10. A composition, comprising a herbicidally effective amount of at least one 3-(heterocyclyl)-substituted benzoylpyrazole of the formula I or an agriculturally useful salt of I as claimed in claim 1 [any of claims 1 to 4] and auxiliaries which are customarily used for formulating crop protection agents.


Claim 11 has been canceled.

12. A method for controlling undesirable vegetation, characterized in that a herbicidally effective amount of at least one 3-(heterocyclyl)-substituted benzoylpyrazole of the formula I or an agriculturally useful salt of I as claimed in claim 1 [any of claims 1 to 4] is allowed to act on the plants, their habitat and/or on seed.

Claim 13 has been canceled.

CURRENT CLAIMS OZ 50555

1. A 3-(heterocyclyl)-substituted benzoylpyrazole of the formula I



where:

X is O, NH or N(C₁-C₆-alkyl);

R¹ is C₁-C₆-alkyl;

R², R³, R⁴, R⁵ are hydrogen, C₁-C₄-alkyl or C₁-C₄-haloalkyl;

R⁶ is halogen, nitro, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio, C₁-C₄-alkylsulfonyl or C₁-C₄-haloalkylsulfonyl;

R⁷ is hydroxyl, C₁-C₆-alkoxy, C₃-C₆-alkenyloxy, C₁-C₆-alkylsulfonyloxy, C₁-C₆-alkylcarbonyloxy, C₁-C₄-(alkylthio)carbonyloxy, phenylsulfonyloxy or phenylcarbonyloxy, where the phenyl radical of the two last-mentioned substituents may be partially or fully halogenated and/or may carry one to three of the following groups:

nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy or C₁-C₄-haloalkoxy;

R^8, R^9 are C_1-C_4 -alkyl;

R^{10} is hydrogen or C_1-C_4 -alkyl;

where the number of the carbon atoms of the radicals R^8 , R^9 and R^{10} together is at most 7,

R^{11} is hydrogen or C_1-C_4 -alkyl;

and its agriculturally useful salts.

2. A 3-(heterocyclyl)-substituted benzoylpyrazole of the formula I as claimed in claim 1 where

X is O;

R^1 is C_1-C_4 -alkyl;

R^6 is C_1-C_4 -alkylthio or C_1-C_4 -alkylsulfonyl.

3. A 3-(heterocyclyl)-substituted benzoylpyrazole of the formula I as claimed in claim 1 where

X is O;

R^1 is C_1-C_4 -alkyl;


R^6 is halogen, nitro, C_1-C_4 -haloalkyl, C_1-C_4 -alkoxy or C_1-C_4 -haloalkoxy.

4. A 3-(heterocyclyl)-substituted benzoylpyrazole of the formula I as claimed in claim 1 where



X is $N(C_1-C_6\text{-alkyl})$.

5. A process for preparing 3-(heterocyclyl)-substituted benzoylpyrazoles of the formula I where R^7 = hydroxyl as claimed in claim 1, which comprises acylating a

pyrazole of the formula II




with an activated benzoic acid III α or a benzoic acid III β ,




where the variables X, R¹ to R⁶ and R⁸ to R¹¹ are as defined in claim 1 and L¹ is a nucleophilically replaceable leaving group and rearranging the acylation product, in the presence or absence of a catalyst, to give the compounds of the formula I where R⁷ = hydroxyl.

6. A process for preparing 3-(heterocyclyl)-substituted benzoylpyrazoles of the formula I where R⁷ = OH as claimed in claim 1, which comprises reacting a pyrazole of the formula II




in which the variables R^8 to R^{11} are as defined in claim 1, or an alkali metal salt thereof, with a 3-(heterocyclyl)benzene derivative of the formula V




where the variables X and R^1 to R^6 are as defined in claim 1 and L^2 is a leaving group in the presence of carbon monoxide, a catalyst and a base.

7. A process for preparing 3-(heterocyclyl)-substituted benzoylpyrazoles of the formula I where $R^7 \neq$ hydroxyl as claimed in claim 1, which comprises reacting a 3-(heterocyclyl)-substituted benzoylpyrazole I where $R^7 =$ hydroxyl



with a compound of the formula VI




where

L^3 is a nucleophilically replaceable leaving group;

R^{7a} is C_1-C_6 -alkyl, C_3-C_6 -alkenyl, C_1-C_6 -alkylsulfonyl, C_1-C_6 -alkylcarbonyl, C_1-C_4 -(alkylthio)carbonyloxy, phenylsulfonyl or phenylcarbonyl, where the phenyl radical of the two last-mentioned substituents may be partially or fully halogenated and/or may carry one to three of the following groups:

nitro, cyano, C_1-C_4 -alkyl, C_1-C_4 -haloalkyl, C_1-C_4 -alkoxy or C_1-C_4 -haloalkoxy.

8. A benzoic acid ester of the formula III




where the variables X, R^1 and R^3 to R^6 are as defined in claim 1 and

R^2 is C_1 – C_4 -haloalkyl; and

L is hydroxyl or a radical that can be removed by hydrolysis.

9. A 3-(heterocyclyl)benzene derivative of the formula V



where the variables X, R^1 and R^3 to R^6 are as defined in claim 1 and

R^2 is C_1 – C_4 -haloalkyl; and

L^2 is a nucleophilically displaceable leaving group.


10. A composition, comprising a herbicidally effective amount of at least one 3-(heterocyclyl)-substituted benzoylpyrazole of the formula I or an agriculturally useful salt of I as claimed in claim 1 and auxiliaries which are customarily used for formulating crop protection agents.

Claim 11 has been canceled.

12. A method for controlling undesirable vegetation, characterized in that a herbicidally effective amount of at least one 3-(heterocyclyl)-substituted benzoylpyrazole of the formula I or an agriculturally useful salt of I as claimed in

3-(Heterocyclyl)-substituted benzoylpyrazoles

The present invention relates to 3-(heterocyclyl)-substituted
5 benzoylpyrazoles of the formula I



where:

- 15 X is O, NH or N(C₁-C₆-alkyl);
- R¹ is C₁-C₆-alkyl;
- 20 R², R³, R⁴, R⁵ are hydrogen, C₁-C₄-alkyl or C₁-C₄-haloalkyl;
- R⁶ is halogen, nitro, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio, C₁-C₄-alkylsulfonyl or C₁-C₄-haloalkylsulfonyl;
- 25 R⁷ is hydroxyl, C₁-C₆-alkoxy, C₃-C₆-alkenyloxy, C₁-C₆-alkylsulfonyloxy, C₁-C₆-alkylcarbonyloxy, C₁-C₄-(alkylthio)carbonyloxy, phenylsulfonyloxy or phenylcarbonyloxy, where the phenyl radical of the two
- 30 last-mentioned substituents may be partially or fully halogenated and/or may carry one to three of the following groups:
nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy or C₁-C₄-haloalkoxy;
- 35 R⁸, R⁹ are C₁-C₄-alkyl;
- R¹⁰ is hydrogen or C₁-C₄-alkyl;
where the number of the carbon atoms of the radicals
- 40 R⁸, R⁹ and R¹⁰ together is at most 7,
- R¹¹ is hydrogen or C₁-C₄-alkyl;

and their agriculturally useful salts.

Moreover, the invention relates to intermediates and processes for preparing compounds of the formula I, to compositions comprising them and to the use of these derivatives or of compositions comprising them for controlling harmful plants.

5

Pyrazol-4-yl benzoyl derivatives are disclosed in the literature, for example in WO 96/26206 and WO 98/31681.

However, the herbicidal properties of the prior-art compounds and
10 their compatibility with crop plants are not entirely satisfactory.

It is an object of the present invention to provide novel, in particular herbicidally active, compounds having improved
15 properties.

We have found that this object is achieved by the 3-(heterocyclyl)-substituted benzoylpyrazoles of the formula I and their herbicidal action.

20

Furthermore, we have found herbicidal compositions which comprise the compounds I and have very good herbicidal action. Moreover, we have found processes for preparing these compositions and methods for controlling undesirable vegetation using the
25 compounds I.

Depending on the substitution pattern, the compounds of the formula I may contain one or more chiral centers, in which case they are present as enantiomers or mixtures of diastereomers. The
30 invention provides both the pure enantiomers or diastereomers and their mixtures.

The compounds of the formula I can also be present in the form of their agriculturally useful salts, the kind of salt usually being
35 immaterial. In general, the salts of those cations or the acid addition salts of those acids are suitable whose cations and anions, respectively, do not adversely affect the herbicidal action of the compounds I.

40 Suitable cations are, in particular, ions of the alkali metals, preferably lithium, sodium and potassium, of the alkaline earth metals, preferably calcium and magnesium, and of the transition metals, preferably manganese, copper, zinc and iron, and also ammonium, where, if desired, one to four hydrogen atoms may be
45 replaced by C₁-C₄-alkyl, hydroxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, hydroxy-C₁-C₄-alkoxy-C₁-C₄-alkyl, phenyl or benzyl, preferably ammonium, dimethylammonium,

3

diisopropylammonium, tetramethylammonium, tetrabutylammonium, 2-(2-hydroxyeth-1-oxy)eth-1-ylammonium, di(2-hydroxyeth-1-yl)ammonium, trimethylbenzylammonium, furthermore phosphonium ions, sulfonium ions, preferably
 5 tri(C₁-C₄-alkyl)sulfonium, and sulfoxonium ions, preferably tri(C₁-C₄-alkyl)sulfoxonium.

Anions of useful acid addition salts are primarily chloride, bromide, fluoride, hydrogen sulfate, sulfate, dihydrogen
 10 phosphate, hydrogen phosphate, nitrate, bicarbonate, carbonate, hexafluorosilicate, hexafluorophosphate, benzoate and the anions of C₁-C₄-alkanoic acids, preferably formate, acetate, propionate and butyrate.

15 The organic molecular moieties mentioned for the substituents R¹-R¹¹ or as radicals on phenyl rings are collective terms for individual enumerations of the individual group members. All hydrocarbon chains, i.e. all alkyl, alkylcarbonyl, haloalkyl, alkoxy, haloalkoxy, alkylcarbonyloxy, (alkylthio)carbonyloxy,
 20 alkylsulfonyloxy, alkylthio, haloalkylthio, alkylsulfonyl, haloalkylsulfonyl, alkenyl and alkenyloxy moieties can be straight-chain or branched. Unless indicated otherwise, halogenated substituents preferably carry one to five identical or different halogen atoms. The term "halogen" represents in each
 25 case fluorine, chlorine, bromine or iodine.

Examples of other meanings are:

- C₁-C₄-alkyl, and the alkyl moieties of C₁-C₄-alkylcarbonyl and
 30 C₁-C₄-alkylcarbonyloxy: for example methyl, ethyl, propyl, 1-methylethyl, butyl, 1-methylpropyl, 2-methylpropyl and 1,1-dimethylethyl;
- C₁-C₆-alkyl, and the alkyl moieties of C₁-C₆-alkylcarbonyl and
 35 C₁-C₆-alkylcarbonyloxy: C₁-C₄-alkyl as mentioned above, and also, for example, pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 2,2-dimethylpropyl, 1-ethylpropyl, hexyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl,
 40 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, 1-ethylbutyl, 2-ethylbutyl, 1,1,2-trimethylpropyl, 1-ethyl-1-methylpropyl and 1-ethyl-3-methylpropyl;
- 45 - C₁-C₄-haloalkyl: a C₁-C₄-alkyl radical as mentioned above which is partially or fully substituted by fluorine, chlorine, bromine and/or iodine, i.e., for example,

- chloromethyl, dichloromethyl, trichloromethyl, fluoromethyl, difluoromethyl, trifluoromethyl, chlorofluoromethyl, dichlorofluoromethyl, chlorodifluoromethyl, bromomethyl, iodomethyl, 1-fluoroethyl, 1-chloroethyl, 1-bromoethyl, 1-iodoethyl, 2-fluoroethyl, 2-chloroethyl, 2-bromoethyl, 2-iodoethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl, 2-chloro-2-fluoroethyl, 2-chloro-2,2-difluoroethyl, 2,2-dichloro-2-fluoroethyl, 2,2,2-trichloroethyl, pentafluoroethyl, 2-fluoropropyl, 3-fluoropropyl, 2,2-difluoropropyl, 2,3-difluoropropyl, 2-chloropropyl, 3-chloropropyl, 2,3-dichloropropyl, 2-bromopropyl, 3-bromopropyl, 3,3,3-trifluoropropyl, 3,3,3-trichloropropyl, 2,2,3,3,3-pentafluoropropyl, heptafluoropropyl, 1-(fluoromethyl)-2-fluoroethyl, 1-(chloromethyl)-2-chloroethyl, 1-(bromomethyl)-2-bromoethyl, 4-fluorobutyl, 4-chlorobutyl, 4-bromobutyl and nonafluorobutyl;
- C₁-C₄-alkoxy: for example methoxy, ethoxy, propoxy, 1-methylethoxy, butoxy, 1-methylpropoxy, 2-methylpropoxy and 1,1-dimethylethoxy;
- C₁-C₆-alkoxy: C₁-C₄-alkoxy as mentioned above, and also, for example, pentoxy, 1-methylbutoxy, 2-methylbutoxy, 3-methylbutoxy, 1,1-dimethylpropoxy, 1,2-dimethylpropoxy, 2,2-dimethylpropoxy, 1-ethylpropoxy, hexoxy, 1-methylpentoxy, 2-methylpentoxy, 3-methylpentoxy, 4-methylpentoxy, 1,1-dimethylbutoxy, 1,2-dimethylbutoxy, 1,3-dimethylbutoxy, 2,2-dimethylbutoxy, 2,3-dimethylbutoxy, 3,3-dimethylbutoxy, 1-ethylbutoxy, 2-ethylbutoxy, 1,1,2-trimethylpropoxy, 1,2,2-trimethylpropoxy, 1-ethyl-1-methylpropoxy and 1-ethyl-2-methylpropoxy;
- C₁-C₄-haloalkoxy: a C₁-C₄-alkoxy radical as mentioned above which is partially or fully substituted by fluorine, chlorine, bromine and/or iodine, i.e., for example, fluoromethoxy, difluoromethoxy, trifluoromethoxy, chlorodifluoromethoxy, bromodifluoromethoxy, 2-fluoroethoxy, 2-chloroethoxy, 2-bromomethoxy, 2-iodoethoxy, 2,2-difluoroethoxy, 2,2,2-trifluoroethoxy, 2-chloro-2-fluoroethoxy, 2-chloro-2,2-difluoroethoxy, 2,2-dichloro-2-fluoroethoxy, 2,2,2-trichloroethoxy, pentafluoroethoxy, 2-fluoropropoxy, 3-fluoropropoxy, 2-chloropropoxy, 3-chloropropoxy, 2-bromopropoxy, 3-bromopropoxy, 2,2-difluoropropoxy, 2,3-difluoropropoxy, 2,3-dichloropropoxy, 3,3,3-trifluoropropoxy, 3,3,3-trichloropropoxy, 2,2,3,3,3-pentafluoropropoxy,

5

heptafluoropropoxy, 1-(fluoromethyl)-2-fluoroethoxy,
1-(chloromethyl)-2-chloroethoxy,
1-(bromomethyl)-2-bromoethoxy, 4-fluorobutoxy,
4-chlorobutoxy, 4-bromobutoxy and nonafluorobutoxy;

5

- C₁-C₄-alkylthio, and the alkylthio moieties of
C₁-C₄-(alkylthio)carbonyloxy: for example methylthio,
ethylthio, propylthio, 1-methylethylthio, butylthio,
1-methylpropylthio, 2-methylpropylthio and
1,1-dimethylethylthio;

10

- C₁-C₄-haloalkylthio: a C₁-C₄-alkylthio radical as mentioned
above which is partially or fully substituted by fluorine,
chlorine, bromine and/or iodine, i.e., for example,
fluoromethylthio, difluoromethylthio, trifluoromethylthio,
chlorodifluoromethylthio, bromodifluoromethylthio,
2-fluoroethylthio, 2-chloroethylthio, 2-bromoethylthio,
2-iodoethylthio, 2,2-difluoroethylthio,
2,2,2-trifluoroethylthio, 2,2,2-trichloroethylthio,
2-chloro-2-fluoroethylthio, 2-chloro-2,2-difluoroethylthio,
2,2-dichloro-2-fluoroethylthio, pentafluoroethylthio,
2-fluoropropylthio, 3-fluoropropylthio, 2-chloropropylthio,
3-chloropropylthio, 2-bromopropylthio, 3-bromopropylthio,
2,2-difluoropropylthio, 2,3-difluoropropylthio,
2,3-dichloropropylthio, 3,3,3-trifluoropropylthio,
3,3,3-trichloropropylthio, 2,2,3,3,3-pentafluoropropylthio,
heptafluoropropylthio, 1-(fluoromethyl)-2-fluoroethylthio,
1-(chloromethyl)-2-chloroethylthio,
1-(bromomethyl)-2-bromoethylthio, 4-fluorobutylthio,
4-chlorobutylthio, 4-bromobutylthio and nonafluorobutylthio;

30

- C₁-C₄-alkylsulfonyl (C₁-C₄-alkyl-S(=O)₂-), and the
alkylsulfonyl moieties of C₁-C₄-alkylsulfonyloxy: for example
methylsulfonyl, ethylsulfonyl, propylsulfonyl,
1-methylethylsulfonyl, butylsulfonyl, 1-methylpropylsulfonyl,
2-methylpropylsulfonyl and 1,1-dimethylethylsulfonyl;

35

- C₁-C₆-alkylsulfonyl, and the alkylsulfonyl moieties of
C₁-C₆-alkylsulfonyloxy: a C₁-C₄-alkylsulfonyl radical as
mentioned above, and also, for example, pentylsulfonyl,
1-methylbutylsulfonyl, 2-methylbutylsulfonyl,
3-methylbutylsulfonyl, 1,1-dimethylpropylsulfonyl,
1,2-dimethylpropylsulfonyl, 2,2-dimethylpropylsulfonyl,
1-ethylpropylsulfonyl, hexylsulfonyl, 1-methylpentylsulfonyl,
2-methylpentylsulfonyl, 3-methylpentylsulfonyl,
4-methylpentylsulfonyl, 1,1-dimethylbutylsulfonyl,
1,2-dimethylbutylsulfonyl, 1,3-dimethylbutylsulfonyl,

45

FBI/DOJ

6

- 2,2-dimethylbutylsulfonyl, 2,3-dimethylbutylsulfonyl,
3,3-dimethylbutylsulfonyl, 1-ethylbutylsulfonyl,
2-ethylbutylsulfonyl, 1,1,2-trimethylpropylsulfonyl,
1,2,2-trimethylpropylsulfonyl, 1-ethyl-1-methylpropylsulfonyl
5 and 1-ethyl-2-methylpropylsulfonyl;
- C₁-C₄-haloalkylsulfonyl: a C₁-C₄-alkylsulfonyl radical as
mentioned above which is partially or fully substituted by
fluorine, chlorine, bromine and/or iodine, i.e., for example,
10 fluoromethylsulfonyl, difluoromethylsulfonyl,
trifluoromethylsulfonyl, chlorodifluoromethylsulfonyl,
bromodifluoromethylsulfonyl, 2-fluoroethylsulfonyl,
2-chloroethylsulfonyl, 2-bromoethylsulfonyl,
2-iodoethylsulfonyl, 2,2-difluoroethylsulfonyl,
15 2,2,2-trifluoroethylsulfonyl, 2-chloro-2-fluoroethylsulfonyl,
2-chloro-2,2-difluoroethylsulfonyl,
2,2-dichloro-2-fluoroethylsulfonyl,
2,2,2-trichloroethylsulfonyl, pentafluoroethylsulfonyl,
2-fluoropropylsulfonyl, 3-fluoropropylsulfonyl,
20 2-chloropropylsulfonyl, 3-chloropropylsulfonyl,
2-bromopropylsulfonyl, 3-bromopropylsulfonyl,
2,2-difluoropropylsulfonyl, 2,3-difluoropropylsulfonyl,
2,3-dichloropropylsulfonyl, 3,3,3-trifluoropropylsulfonyl,
3,3,3-trichloropropylsulfonyl,
25 2,2,3,3,3-pentafluoropropylsulfonyl,
heptafluoropropylsulfonyl,
1-(fluoromethyl)-2-fluoroethylsulfonyl, 1-(chloromethyl)-2-
chloroethylsulfonyl, 1-(bromomethyl)-2-bromoethylsulfonyl,
4-fluorobutylsulfonyl, 4-chlorobutylsulfonyl,
30 4-bromobutylsulfonyl and nonafluorobutylsulfonyl;
- C₃-C₆-alkenyloxy: for example prop-1-en-1-yloxy,
prop-2-en-1-yloxy, 1-methylethenyloxy, buten-1-yloxy,
buten-2-yloxy, buten-3-yloxy, 1-methylprop-1-en-1-yloxy,
35 2-methylprop-1-en-1-yloxy, 1-methylprop-2-en-1-yloxy,
2-methylprop-2-en-1-yloxy, penten-1-yloxy, penten-2-yloxy,
penten-3-yloxy, penten-4-yloxy, 1-methylbut-1-en-1-yloxy,
2-methylbut-1-en-1-yloxy, 3-methylbut-1-en-1-yloxy,
1-methylbut-2-en-1-yloxy, 2-methylbut-2-en-1-yloxy,
40 3-methylbut-2-en-1-yloxy, 1-methylbut-3-en-1-yloxy,
2-methylbut-3-en-1-yloxy, 3-methylbut-3-en-1-yloxy,
1,1-dimethylprop-2-en-1-yloxy, 1,2-dimethylprop-1-en-1-yloxy,
1,2-dimethylprop-2-en-1-yloxy, 1-ethylprop-1-en-2-yloxy,
1-ethylprop-2-en-1-yloxy, hex-1-en-1-yloxy, hex-2-en-1-yloxy,
45 hex-3-en-1-yloxy, hex-4-en-1-yloxy, Hex-5-en-1-yloxy,
1-methylpent-1-en-1-yloxy, 2-methylpent-1-en-1-yloxy,
3-methylpent-1-en-1-yloxy, 4-methylpent-1-en-1-yloxy,

- 1-methylpent-2-en-1-yloxy, 2-methylpent-2-en-1-yloxy,
 3-methylpent-2-en-1-yloxy, 4-methylpent-2-en-1-yloxy,
 1-methylpent-3-en-1-yloxy, 2-methylpent-3-en-1-yloxy,
 3-methylpent-3-en-1-yloxy, 4-methylpent-3-en-1-yloxy,
 5 1-methylpent-4-en-1-yloxy, 2-methylpent-4-en-1-yloxy,
 3-methylpent-4-en-1-yloxy, 4-methylpent-4-en-1-yloxy,
 1,1-dimethylbut-2-en-1-yloxy, 1,1-dimethylbut-3-en-1-yloxy,
 1,2-dimethylbut-1-en-1-yloxy, 1,2-dimethylbut-2-en-1-yloxy,
 1,2-dimethylbut-3-en-1-yloxy, 1,3-dimethylbut-1-en-1-yloxy,
 10 1,3-dimethylbut-2-en-1-yloxy, 1,3-dimethylbut-3-en-1-yloxy,
 2,2-dimethylbut-3-en-1-yloxy, 2,3-dimethylbut-1-en-1-yloxy,
 2,3-dimethylbut-2-en-1-yloxy, 2,3-dimethylbut-3-en-1-yloxy,
 3,3-dimethylbut-1-en-1-yloxy, 3,3-dimethylbut-2-en-1-yloxy,
 1-ethylbut-1-en-1-yloxy, 1-ethylbut-2-en-1-yloxy,
 15 1-ethylbut-3-en-1-yloxy, 2-ethylbut-1-en-1-yloxy,
 2-ethylbut-2-en-1-yloxy, 2-ethylbut-3-en-1-yloxy,
 1,1,2-trimethylprop-2-en-1-yloxy,
 1-ethyl-1-methylprop-2-en-1-yloxy,
 1-ethyl-2-methylprop-1-en-1-yloxy and
 20 1-ethyl-2-methylprop-2-en-1-
 yloxy;
- C₃-C₆-alkenyl: prop-1-en-1-yl, prop-2-en-1-yl,
 1-methylethenyl, buten-1-yl, buten-2-yl, buten-3-yl,
 25 1-methylprop-1-en-1-yl, 2-methylprop-1-en-1-yl,
 1-methylprop-2-en-1-yl, 2-methylprop-2-en-1-yl, penten-1-yl,
 penten-2-yl, penten-3-yl, penten-4-yl, 1-methylbut-1-en-1-yl,
 2-methylbut-1-en-1-yl, 3-methylbut-1-en-1-yl,
 1-methylbut-2-en-1-yl, 2-methylbut-2-en-1-yl,
 30 3-methylbut-2-en-1-yl, 1-methylbut-3-en-1-yl,
 2-methylbut-3-en-1-yl, 3-methylbut-3-en-1-yl,
 1,1-dimethylprop-2-en-1-yl, 1,2-dimethylprop-1-en-1-yl,
 1,2-dimethylprop-2-en-1-yl, 1-ethylprop-1-en-2-yl,
 1-ethylprop-2-en-1-yl, hex-1-en-1-yl, hex-2-en-1-yl,
 35 hex-3-en-1-yl, hex-4-en-1-yl, hex-5-en-1-yl,
 1-methylpent-1-en-1-yl, 2-methylpent-1-en-1-yl,
 3-methylpent-1-en-1-yl, 4-methylpent-1-en-1-yl,
 1-methylpent-2-en-1-yl, 2-methylpent-2-en-1-yl,
 3-methylpent-2-en-1-yl, 4-methylpent-2-en-1-yl,
 40 1-methylpent-3-en-1-yl, 2-methylpent-3-en-1-yl,
 3-methylpent-3-en-1-yl, 4-methylpent-3-en-1-yl,
 1-methylpent-4-en-1-yl, 2-methylpent-4-en-1-yl,
 3-methylpent-4-en-1-yl, 4-methylpent-4-en-1-yl,
 1,1-dimethylbut-2-en-1-yl, 1,1-dimethylbut-3-en-1-yl,
 45 1,2-dimethylbut-1-en-1-yl, 1,2-dimethylbut-2-en-1-yl,
 1,2-dimethylbut-3-en-1-yl, 1,3-dimethylbut-1-en-1-yl,
 1,3-dimethylbut-2-en-1-yl, 1,3-dimethylbut-3-en-1-yl,

- 2,2-dimethylbut-3-en-1-yl, 2,3-dimethylbut-1-en-1-yl,
 2,3-dimethylbut-2-en-1-yl, 2,3-dimethylbut-3-en-1-yl,
 3,3-dimethylbut-1-en-1-yl, 3,3-dimethylbut-2-en-1-yl,
 1-ethylbut-1-en-1-yl, 1-ethylbut-2-en-1-yl,
 5 1-ethylbut-3-en-1-yl, 2-ethylbut-1-en-1-yl,
 2-ethylbut-2-en-1-yl, 2-ethylbut-3-en-1-yl,
 1,1,2-trimethylprop-2-en-1-yl,
 1-ethyl-1-methylprop-2-en-1-yl,
 1-ethyl-2-methylprop-1-en-1-yl and
 10 1-ethyl-2-methylprop-2-en-1-yl.

The phenyl rings are preferably unsubstituted or carry one to three halogen atoms and/or one nitro group, one cyano group, one or two methyl, trifluoromethyl, methoxy or trifluoromethoxy
 15 groups.

Emphasis is given to those 3-(heterocyclyl)-substituted benzoylpyrazoles of the formula I where

- 20 R⁷ is hydroxyl, C₁-C₆-alkoxy, C₃-C₆-alkenyloxy, C₁-C₆-alkylsulfonyloxy, C₁-C₆-alkylcarbonyloxy, phenylsulfonyloxy or phenylcarbonyloxy, where the phenyl radical of the two last-mentioned substituents may be partially or fully
 25 halogenated and/or may carry one to three of the following groups: nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy or C₁-C₄-haloalkoxy.
- Preference is given to the 3-(heterocyclyl)-substituted benzoylpyrazoles of the formula I where:
 30 X is O;
 R¹ is C₁-C₄-alkyl;
 particularly preferably methyl or ethyl;
 35 with particular preference methyl;
 R², R³, R⁴, R⁵ are hydrogen, C₁-C₄-alkyl or C₁-C₄-haloalkyl;
 particularly preferably hydrogen, methyl, ethyl,
 40 propyl, 1-methylethyl, fluoromethyl or chloromethyl;
 with particular preference hydrogen, methyl, ethyl or chloromethyl;
 R⁶ is C₁-C₄-alkylthio or C₁-C₄-alkylsulfonyl;
 45 particularly preferably methylthio, ethylthio or 1-methyl-1-ethylthio, methylsulfonyl, ethylsulfonyl, 1-methylethylsulfonyl or

propylsulfonyl;
with particular preference methylsulfonyl,
ethylsulfonyl, 1-methylethylsulfonyl or
propylsulfonyl;

5

R⁷ is hydroxyl, C₁-C₆-alkoxy, C₃-C₆-alkenyloxy,
C₁-C₆-alkylsulfonyloxy, C₁-C₆-alkylcarbonyloxy,
C₁-C₄-(alkylthio)carbonyloxy, phenylsulfonyloxy or
phenylcarbonyloxy, where the phenyl radical of the
two last-mentioned substituents may be partially
or fully halogenated and/or may carry one to three
of the following groups:

10

nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl,
C₁-C₄-alkoxy or C₁-C₄-haloalkoxy;

15

particularly preferably hydroxyl, C₁-C₄-alkoxy,
C₃-C₆-alkenyloxy, C₁-C₄-alkylsulfonyloxy,
C₁-C₄-alkylcarbonyloxy, phenylsulfonyloxy or
phenylcarbonyloxy, where the phenyl radical of the
two last-mentioned substituents may be partially
or fully halogenated and/or may carry one to three
of the following groups:

20

nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl,
C₁-C₄-alkoxy or C₁-C₄-haloalkoxy;

25

R⁸, R⁹

are C₁-C₄-alkyl;
particularly preferably methyl, ethyl, propyl,
1-methyl-1-ethyl, butyl, 1-methyl-1-propyl and
2-methyl-1-propyl;

30

R¹⁰

is hydrogen or C₁-C₄-alkyl;
particularly preferably C₁-C₄-alkyl;
with particular preference methyl, ethyl or
propyl;

35

R¹¹

is hydrogen or C₁-C₄-alkyl;
particularly preferably hydrogen or methyl.

Particular preference is given to the
3-(heterocyclyl)-substituted benzoylpyrazoles of the formula
I where

40

X is O;

R¹

45

is C₁-C₄-alkyl;
particularly preferably methyl or ethyl;
with particular preference methyl;

10

R⁶ is C₁-C₄-alkylsulfonyl;
particularly preferably methylsulfonyl,
ethylsulfonyl, 1-methyl-1-ethylsulfonyl or
propylsulfonyl;

5

R⁷ is hydroxyl, C₁-C₆-alkylsulfonyloxy,
C₁-C₆-alkylcarbonyloxy, phenylsulfonyloxy or
phenylcarbonyloxy, where the phenyl radical of the
two last-mentioned substituents may be partially
or fully halogenated and/or may carry one to three
of the following groups:
nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl,
C₁-C₄-alkoxy or C₁-C₄-haloalkoxy;
particularly preferably hydroxyl;

10

15

R⁸, R⁹ are C₁-C₄-alkyl;

R¹⁰ is hydrogen or C₁-C₄-alkyl.

20

Very particular preference is given to the
3-(heterocyclyl)-substituted benzoylpyrazoles of the formula
I where

R⁸ is C₂-C₄-alkyl, for example ethyl,
1-methyl-1-ethyl, propyl or butyl;

25

R⁹ is C₁-C₄-alkyl, for example methyl or ethyl;

R¹⁰ is hydrogen or C₁-C₄-alkyl, for example methyl or
ethyl.

30

Very particular preference is also given to the
3-(heterocyclyl)-substituted benzoylpyrazoles of the formula
I where

35

R⁸ is methyl;

R⁹ is C₁-C₄-alkyl, for example methyl, ethyl, propyl
or butyl;

40

R¹⁰ is C₁-C₄-alkyl, for example methyl or ethyl.

Very particular preference is also given to the
3-(heterocyclyl)-substituted benzoylpyrazoles of the formula
I where

45

- R^8, R^9 are methyl;
- R^{10} is hydrogen.
- 5 • Preference is also given to the 3-(heterocyclyl)-substituted benzoylpyrazoles of the formula I where:
- X is O
- 10 R^1 is C_1 - C_4 -alkyl;
particularly preferably methyl or ethyl;
with particular preference methyl;
- 15 R^2, R^3, R^4, R^5 are hydrogen, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl;
particularly preferably hydrogen, methyl, ethyl,
propyl, 1-methyl-1-ethyl, chloromethyl or
fluoromethyl;
with particular preference hydrogen, methyl, ethyl
or chloromethyl;
- 20 R^6 is halogen, nitro, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy or
 C_1 - C_4 -haloalkoxy;
particularly preferably halogen, such as chlorine
or bromine, nitro, C_1 - C_2 -haloalkyl, such as
- 25 difluoromethyl or trifluoromethyl, C_1 - C_2 -alkoxy or
 C_1 - C_2 -haloalkoxy, such as difluoromethoxy,
chlorodifluoromethoxy or trifluoromethoxy;
- 30 R^7 is hydroxyl, C_1 - C_6 -alkoxy, C_3 - C_6 -alkenyloxy,
 C_1 - C_6 -alkylsulfonyloxy, C_1 - C_6 -alkylcarbonyloxy,
 C_1 - C_4 -(alkylthio)carbonyloxy, phenylsulfonyloxy or
phenylcarbonyloxy, where the phenyl radical of the
two last-mentioned substituents may be partially
or fully halogenated and/or may carry one to three
- 35 of the following groups:
nitro, cyano, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl,
 C_1 - C_4 -alkoxy or C_1 - C_4 -haloalkoxy;
particularly preferably hydroxyl, C_1 - C_4 -alkoxy,
 C_3 - C_6 -alkenyloxy, C_1 - C_4 -alkylsulfonyloxy,
- 40 C_1 - C_4 -alkylcarbonyloxy, phenylsulfonyloxy or
phenylcarbonyloxy, where the phenyl radical of the
two last-mentioned substituents may be partially
or fully halogenated and/or may carry one to three
of the following groups:
- 45 nitro, cyano, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl,
 C_1 - C_4 -alkoxy or C_1 - C_4 -haloalkoxy;

12

R^8, R^9 are C_1-C_4 -alkyl;
particularly preferably methyl, ethyl, propyl,
1-methyl-1-ethyl, butyl, 1-methyl-1-propyl and
2-methyl-1-propyl;

5

R^{10} is hydrogen or C_1-C_4 -alkyl;
particularly preferably C_1-C_4 -alkyl;
with particular preference methyl, ethyl or
propyl;

10

R^{11} is hydrogen or C_1-C_4 -alkyl;
particularly preferably hydrogen or methyl.

Particular preference is given to the
15 3-(heterocyclyl)-substituted benzoylpyrazoles of the formula
I where

X is O;

20

R^1 is C_1-C_4 -alkyl;
particularly preferably methyl or ethyl;
with particular preference methyl;

25

R^6 is halogen, nitro, C_1-C_4 -haloalkyl, C_1-C_4 -alkoxy or
 C_1-C_4 -haloalkoxy;
particularly preferably halogen, such as chlorine
or bromine, nitro, C_1-C_2 -haloalkyl, such as
difluoromethyl or trifluoromethyl, C_1-C_2 -alkoxy or
 C_1-C_2 -haloalkoxy, such as difluoromethoxy;

30

R^7 is hydroxyl, C_1-C_6 -alkylsulfonyloxy,
 C_1-C_6 -alkylcarbonyloxy, phenylsulfonyloxy or
phenylcarbonyloxy, where the phenyl radical of the
two last-mentioned substituents may be partially
35 or fully halogenated and/or may carry one to three
of the following groups:
nitro, cyano, C_1-C_4 -alkyl, C_1-C_4 -haloalkyl,
 C_1-C_4 -alkoxy or C_1-C_4 -haloalkoxy;
particularly preferably hydroxyl;

40

R^8, R^9 are C_1-C_4 -alkyl;

R^{10} is hydrogen or C_1-C_4 -alkyl.

45

5 R⁸ is C₂-C₄-alkyl, for example ethyl,
1-methyl-1-ethyl, propyl or butyl;

10 R^{10} is hydrogen or C₁-C₄-alkyl, for example methyl or ethyl.

15

20 R⁹ is C₁-C₄-alkyl, for example methyl, ethyl, propyl
or butyl;

20

25

30 R^{10} is hydrogen.

35 X is N(C₁-C₆-alkyl);
particularly preferably N-methyl, N-ethyl,
N-(1-methyl-1-ethyl) or N-propyl;

40

45

with particular preference hydrogen, methyl, ethyl or chloromethyl;

5 R⁶ is halogen, nitro, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio, C₁-C₄-alkylsulfonyl or C₁-C₄-haloalkylsulfonyl; particularly preferably halogen, such as fluorine, chlorine or bromine, nitro, C₁-C₄-haloalkyl, such as difluoromethyl or trifluoromethyl, C₁-C₄-alkoxy, such as methoxy or ethoxy, C₁-C₄-haloalkoxy, such as difluoromethoxy, chlorodifluoromethoxy or trifluoromethoxy, C₁-C₄-alkylthio, such as methylthio or ethylthio, or C₁-C₄-alkylsulfonyl, such as methylsulfonyl, ethylsulfonyl, 1-methyl-1-ethylsulfonyl or propylsulfonyl;

20 R⁷ is hydroxyl, C₁-C₆-alkoxy, C₃-C₆-alkenyloxy, C₁-C₆-alkylsulfonyloxy, C₁-C₆-alkylcarbonyloxy, C₁-C₄-(alkylthio)carbonyloxy, phenylsulfonyloxy or phenylcarbonyloxy, where the phenyl radical of the two last-mentioned substituents may be partially or fully halogenated and/or may carry one to three of the following groups:
25 nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy or C₁-C₄-haloalkoxy; particularly preferably hydroxyl, C₁-C₄-alkoxy, C₃-C₆-alkenyloxy, C₁-C₄-alkylsulfonyloxy, C₁-C₄-alkylcarbonyloxy, phenylsulfonyloxy or
30 phenylcarbonyloxy, where the phenyl radical of the two last-mentioned substituents may be partially or fully halogenated and/or may carry one to three of the following groups:
35 nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy or C₁-C₄-haloalkoxy;

40 R⁸, R⁹ are C₁-C₄-alkyl; particularly preferably methyl, ethyl, propyl, 1-methyl-1-ethyl, butyl, 1-methyl-1-propyl and 2-methyl-1-propyl;

45 R¹⁰ is hydrogen or C₁-C₄-alkyl; particularly preferably C₁-C₄-alkyl; with particular preference methyl, ethyl or propyl;

15

R¹¹ is hydrogen or C₁-C₄-alkyl;
particularly preferably hydrogen or methyl.

5 Particular preference is given to the
3-heterocyclyl-substituted benzoylpyrazoles of the formula I
where

R⁷ is hydroxyl, C₁-C₆-alkylsulfonyloxy,
10 C₁-C₆-alkylcarbonyloxy, phenylsulfonyloxy or
phenylcarbonyloxy, where the phenyl radical of the
two last-mentioned substituents may be partially
or fully halogenated and/or may carry one to three
of the following groups:
15 nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl,
C₁-C₄-alkoxy or C₁-C₄-haloalkoxy;
particularly preferably hydroxyl.

• Preference is likewise given to the 3-(heterocyclyl)-
20 substituted benzoylpyrazoles of the formula I in which the
variables are as defined below:

X is O;

25 R¹ is C₁-C₄-alkyl;
particularly preferably methyl or ethyl;

R², R³, R⁴, R⁵ are hydrogen, C₁-C₄-alkyl or C₁-C₄-haloalkyl;
particularly preferably hydrogen or C₁-C₄-haloalkyl;

30 R⁶ is C₁-C₄-alkylsulfonyl;
particularly preferably methylsulfonyl;

35 R⁷ is hydroxyl, C₁-C₆-alkoxy, C₁-C₆-alkylcarbonyloxy,
C₁-C₄-(alkylthio)carbonyloxy, phenylsulfonyloxy or
phenylcarbonyloxy, where the phenyl radical of the
two last-mentioned substituents may be partially or
fully halogenated and/or may carry one to three of
the following groups:
40 nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl,
C₁-C₄-alkoxy or C₁-C₄-haloalkoxy;
particularly preferably hydroxyl, C₁-C₄-alkoxy or
phenylcarbonyloxy, where the phenyl radical may be
[lacuna] or fully halogenated and/or may carry one
45 to three C₁-C₄-haloalkyl radicals;

R⁸, R⁹ are C₁-C₄-alkyl;

R^{10} is hydrogen or C_1 - C_4 -alkyl;


R^{11} is hydrogen or C_1 - C_4 -alkyl.

5

Very particular preference is given to the compounds of the formula Ia1 (\equiv I where $R^1, R^8, R^9 = CH_3$; $R^{10}, R^{11} = H$), in particular to the compounds Ia1.1 to Ia1.300 of Table 1, where the radical definitions X and R^1 to R^{11} are of particular importance for the compounds according to the invention, not only in combination with one another, but in each case also on their own.

10

15



Ia1

20

Table 1:

25

30

35

40

45

No.	X	R^2	R^3	R^4	R^5	R^6	R^7
Ia1.1	O	H	H	H	H	SCH ₃	OH
Ia1.2	O	H	H	H	H	SCH ₂ CH ₃	OH
Ia1.3	O	H	H	H	H	SO ₂ CH ₃	OH
Ia1.4	O	H	H	H	H	SO ₂ CH ₂ CH ₃	OH
Ia1.5	O	H	H	H	H	SO ₂ CH(CH ₃) ₂	OH
Ia1.6	O	H	H	H	H	SO ₂ (CH ₂) ₂ CH ₃	OH
Ia1.7	O	H	H	H	H	Cl	OH
Ia1.8	O	H	H	H	H	Br	OH
Ia1.9	O	H	H	H	H	NO ₂	OH
Ia1.10	O	H	H	H	H	CHF ₂	OH
Ia1.11	O	H	H	H	H	CF ₃	OH
Ia1.12	O	H	H	H	H	OCH ₃	OH
Ia1.13	O	H	H	H	H	OCH ₂ CH ₃	OH
Ia1.14	O	H	H	H	H	OCHF ₂	OH
Ia1.15	O	H	H	H	H	OCF ₃	OH
Ia1.16	O	CH ₃	H	H	H	SCH ₃	OH
Ia1.17	O	CH ₃	H	H	H	SCH ₂ CH ₃	OH
Ia1.18	O	CH ₃	H	H	H	SO ₂ CH ₃	OH
Ia1.19	O	CH ₃	H	H	H	SO ₂ CH ₂ CH ₃	OH
Ia1.20	O	CH ₃	H	H	H	SO ₂ CH(CH ₃) ₂	OH
Ia1.21	O	CH ₃	H	H	H	SO ₂ (CH ₂) ₂ CH ₃	OH

	No.	X	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷
5	Ia1.22	O	CH ₃	H	H	H	Cl	OH
	Ia1.23	O	CH ₃	H	H	H	Br	OH
	Ia1.24	O	CH ₃	H	H	H	NO ₂	OH
	Ia1.25	O	CH ₃	H	H	H	CHF ₂	OH
	Ia1.26	O	CH ₃	H	H	H	CF ₃	OH
10	Ia1.27	O	CH ₃	H	H	H	OCH ₃	OH
	Ia1.28	O	CH ₃	H	H	H	OCH ₂ CH ₃	OH
	Ia1.29	O	CH ₃	H	H	H	OCHF ₂	OH
	Ia1.30	O	CH ₃	H	H	H	OCF ₃	OH
	Ia1.31	O	H	H	CH ₃	H	SCH ₃	OH
15	Ia1.32	O	H	H	CH ₃	H	SCH ₂ CH ₃	OH
	Ia1.33	O	H	H	CH ₃	H	SO ₂ CH ₃	OH
	Ia1.34	O	H	H	CH ₃	H	SO ₂ CH ₂ CH ₃	OH
	Ia1.35	O	H	H	CH ₃	H	SO ₂ CH(CH ₃) ₂	OH
	Ia1.36	O	H	H	CH ₃	H	SO ₂ (CH ₂) ₂ CH ₃	OH
20	Ia1.37	O	H	H	CH ₃	H	Cl	OH
	Ia1.38	O	H	H	CH ₃	H	Br	OH
	Ia1.39	O	H	H	CH ₃	H	NO ₂	OH
	Ia1.40	O	H	H	CH ₃	H	CHF ₂	OH
	Ia1.41	O	H	H	CH ₃	H	CF ₃	OH
25	Ia1.42	O	H	H	CH ₃	H	OCH ₃	OH
	Ia1.43	O	H	H	CH ₃	H	OCH ₂ CH ₃	OH
	Ia1.44	O	H	H	CH ₃	H	OCHF ₂	OH
	Ia1.45	O	H	H	CH ₃	H	OCF ₃	OH
	Ia1.46	O	CH ₃	CH ₃	H	H	SCH ₃	OH
30	Ia1.47	O	CH ₃	CH ₃	H	H	SCH ₂ CH ₃	OH
	Ia1.48	O	CH ₃	CH ₃	H	H	SO ₂ CH ₃	OH
	Ia1.49	O	CH ₃	CH ₃	H	H	SO ₂ CH ₂ CH ₃	OH
	Ia1.50	O	CH ₃	CH ₃	H	H	SO ₂ CH(CH ₃) ₂	OH
	Ia1.51	O	CH ₃	CH ₃	H	H	SO ₂ (CH ₂) ₂ CH ₃	OH
35	Ia1.52	O	CH ₃	CH ₃	H	H	Cl	OH
	Ia1.53	O	CH ₃	CH ₃	H	H	Br	OH
	Ia1.54	O	CH ₃	CH ₃	H	H	NO ₂	OH
	Ia1.55	O	CH ₃	CH ₃	H	H	CHF ₂	OH
	Ia1.56	O	CH ₃	CH ₃	H	H	CF ₃	OH
40	Ia1.57	O	CH ₃	CH ₃	H	H	OCH ₃	OH
	Ia1.58	O	CH ₃	CH ₃	H	H	OCH ₂ CH ₃	OH
	Ia1.59	O	CH ₃	CH ₃	H	H	OCHF ₂	OH
	Ia1.60	O	CH ₃	CH ₃	H	H	OCF ₃	OH

	No.	X	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷
5	Ial.61	O	CH ₃	H	CH ₃	H	SCH ₃	OH
	Ial.62	O	CH ₃	H	CH ₃	H	SCH ₂ CH ₃	OH
	Ial.63	O	CH ₃	H	CH ₃	H	SO ₂ CH ₃	OH
	Ial.64	O	CH ₃	H	CH ₃	H	SO ₂ CH ₂ CH ₃	OH
	Ial.65	O	CH ₃	H	CH ₃	H	SO ₂ CH(CH ₃) ₂	OH
10	Ial.66	O	CH ₃	H	CH ₃	H	SO ₂ (CH ₂) ₂ CH ₃	OH
	Ial.67	O	CH ₃	H	CH ₃	H	Cl	OH
	Ial.68	O	CH ₃	H	CH ₃	H	Br	OH
	Ial.69	O	CH ₃	H	CH ₃	H	NO ₂	OH
	Ial.70	O	CH ₃	H	CH ₃	H	CHF ₂	OH
15	Ial.71	O	CH ₃	H	CH ₃	H	CF ₃	OH
	Ial.72	O	CH ₃	H	CH ₃	H	OCH ₃	OH
	Ial.73	O	CH ₃	H	CH ₃	H	OCH ₂ CH ₃	OH
	Ial.74	O	CH ₃	H	CH ₃	H	OCHF ₂	OH
	Ial.75	O	CH ₃	H	CH ₃	H	OCF ₃	OH
20	Ial.76	O	H	H	CH ₃	CH ₃	SCH ₃	OH
	Ial.77	O	H	H	CH ₃	CH ₃	SCH ₂ CH ₃	OH
	Ial.78	O	H	H	CH ₃	CH ₃	SO ₂ CH ₃	OH
	Ial.79	O	H	H	CH ₃	CH ₃	SO ₂ CH ₂ CH ₃	OH
	Ial.80	O	H	H	CH ₃	CH ₃	SO ₂ CH(CH ₃) ₂	OH
25	Ial.81	O	H	H	CH ₃	CH ₃	SO ₂ (CH ₂) ₂ CH ₃	OH
	Ial.82	O	H	H	CH ₃	CH ₃	Cl	OH
	Ial.83	O	H	H	CH ₃	CH ₃	Br	OH
	Ial.84	O	H	H	CH ₃	CH ₃	NO ₂	OH
	Ial.85	O	H	H	CH ₃	CH ₃	CHF ₂	OH
30	Ial.86	O	H	H	CH ₃	CH ₃	CF ₃	OH
	Ial.87	O	H	H	CH ₃	CH ₃	OCH ₃	OH
	Ial.88	O	H	H	CH ₃	CH ₃	OCH ₂ CH ₃	OH
	Ial.89	O	H	H	CH ₃	CH ₃	OCHF ₂	OH
	Ial.90	O	H	H	CH ₃	CH ₃	OCF ₃	OH
35	Ial.91	O	CH ₃	CH ₃	CH ₃	H	SCH ₃	OH
	Ial.92	O	CH ₃	CH ₃	CH ₃	H	SCH ₂ CH ₃	OH
	Ial.93	O	CH ₃	CH ₃	CH ₃	H	SO ₂ CH ₃	OH
	Ial.94	O	CH ₃	CH ₃	CH ₃	H	SO ₂ CH ₂ CH ₃	OH
	Ial.95	O	CH ₃	CH ₃	CH ₃	H	SO ₂ CH(CH ₃) ₂	OH
40	Ial.96	O	CH ₃	CH ₃	CH ₃	H	SO ₂ (CH ₂) ₂ CH ₃	OH
	Ial.97	O	CH ₃	CH ₃	CH ₃	H	Cl	OH
	Ial.98	O	CH ₃	CH ₃	CH ₃	H	Br	OH
	Ial.99	O	CH ₃	CH ₃	CH ₃	H	NO ₂	OH

	No.	X	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷
5	Ia1.100	O	CH ₃	CH ₃	CH ₃	H	CHF ₂	OH
	Ia1.101	O	CH ₃	CH ₃	CH ₃	H	CF ₃	OH
	Ia1.102	O	CH ₃	CH ₃	CH ₃	H	OCH ₃	OH
	Ia1.103	O	CH ₃	CH ₃	CH ₃	H	OCH ₂ CH ₃	OH
	Ia1.104	O	CH ₃	CH ₃	CH ₃	H	OCHF ₂	OH
10	Ia1.105	O	CH ₃	CH ₃	CH ₃	H	OCF ₃	OH
	Ia1.106	O	CH ₃	H	CH ₃	CH ₃	SCH ₃	OH
	Ia1.107	O	CH ₃	H	CH ₃	CH ₃	SCH ₂ CH ₃	OH
	Ia1.108	O	CH ₃	H	CH ₃	CH ₃	SO ₂ CH ₃	OH
	Ia1.109	O	CH ₃	H	CH ₃	CH ₃	SO ₂ CH ₂ CH ₃	OH
15	Ia1.110	O	CH ₃	H	CH ₃	CH ₃	SO ₂ CH(CH ₃) ₂	OH
	Ia1.111	O	CH ₃	H	CH ₃	CH ₃	SO ₂ (CH ₂) ₂ CH ₃	OH
	Ia1.112	O	CH ₃	H	CH ₃	CH ₃	Cl	OH
	Ia1.113	O	CH ₃	H	CH ₃	CH ₃	Br	OH
	Ia1.114	O	CH ₃	H	CH ₃	CH ₃	NO ₂	OH
20	Ia1.115	O	CH ₃	H	CH ₃	CH ₃	CHF ₂	OH
	Ia1.116	O	CH ₃	H	CH ₃	CH ₃	CF ₃	OH
	Ia1.117	O	CH ₃	H	CH ₃	CH ₃	OCH ₃	OH
	Ia1.118	O	CH ₃	H	CH ₃	CH ₃	OCH ₂ CH ₃	OH
	Ia1.119	O	CH ₃	H	CH ₃	CH ₃	OCHF ₂	OH
25	Ia1.120	O	CH ₃	H	CH ₃	CH ₃	OCF ₃	OH
	Ia1.121	O	CH ₃	CH ₃	CH ₃	CH ₃	SCH ₃	OH
	Ia1.122	O	CH ₃	CH ₃	CH ₃	CH ₃	SCH ₂ CH ₃	OH
	Ia1.123	O	CH ₃	CH ₃	CH ₃	CH ₃	SO ₂ CH ₃	OH
	Ia1.124	O	CH ₃	CH ₃	CH ₃	CH ₃	SO ₂ CH ₂ CH ₃	OH
30	Ia1.125	O	CH ₃	CH ₃	CH ₃	CH ₃	SO ₂ CH(CH ₃) ₂	OH
	Ia1.126	O	CH ₃	CH ₃	CH ₃	CH ₃	SO ₂ (CH ₂) ₂ CH ₃	OH
	Ia1.127	O	CH ₃	CH ₃	CH ₃	CH ₃	Cl	OH
	Ia1.128	O	CH ₃	CH ₃	CH ₃	CH ₃	Br	OH
	Ia1.129	O	CH ₃	CH ₃	CH ₃	CH ₃	NO ₂	OH
35	Ia1.130	O	CH ₃	CH ₃	CH ₃	CH ₃	CHF ₂	OH
	Ia1.131	O	CH ₃	CH ₃	CH ₃	CH ₃	CF ₃	OH
	Ia1.132	O	CH ₃	CH ₃	CH ₃	CH ₃	OCH ₃	OH
	Ia1.133	O	CH ₃	CH ₃	CH ₃	CH ₃	OCH ₂ CH ₃	OH
	Ia1.134	O	CH ₃	CH ₃	CH ₃	CH ₃	OCHF ₂	OH
40	Ia1.135	O	CH ₃	CH ₃	CH ₃	CH ₃	OCF ₃	OH
	Ia1.136	O	CH ₂ Cl	H	H	H	SCH ₃	OH
	Ia1.137	O	CH ₂ Cl	H	H	H	SCH ₂ CH ₃	OH
	Ia1.138	O	CH ₂ Cl	H	H	H	SO ₂ CH ₃	OH

	No.	X	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷
5	Ial.139	O	CH ₂ Cl	H	H	H	SO ₂ CH ₂ CH ₃	OH
	Ial.140	O	CH ₂ Cl	H	H	H	SO ₂ CH(CH ₃) ₂	OH
	Ial.141	O	CH ₂ Cl	H	H	H	SO ₂ (CH ₂) ₂ CH ₃	OH
	Ial.142	O	CH ₂ Cl	H	H	H	Cl	OH
	Ial.143	O	CH ₂ Cl	H	H	H	Br	OH
10	Ial.144	O	CH ₂ Cl	H	H	H	NO ₂	OH
	Ial.145	O	CH ₂ Cl	H	H	H	CHF ₂	OH
	Ial.146	O	CH ₂ Cl	H	H	H	CF ₃	OH
	Ial.147	O	CH ₂ Cl	H	H	H	OCH ₃	OH
	Ial.148	O	CH ₂ Cl	H	H	H	OCH ₂ CH ₃	OH
15	Ial.149	O	CH ₂ Cl	H	H	H	OCHF ₂	OH
	Ial.150	O	CH ₂ Cl	H	H	H	OCF ₃	OH
	Ial.151	NCH ₃	H	H	H	H	SCH ₃	OH
	Ial.152	NCH ₃	H	H	H	H	SCH ₂ CH ₃	OH
	Ial.153	NCH ₃	H	H	H	H	SO ₂ CH ₃	OH
20	Ial.154	NCH ₃	H	H	H	H	SO ₂ CH ₂ CH ₃	OH
	Ial.155	NCH ₃	H	H	H	H	SO ₂ CH(CH ₃) ₂	OH
	Ial.156	NCH ₃	H	H	H	H	SO ₂ (CH ₂) ₂ CH ₃	OH
	Ial.157	NCH ₃	H	H	H	H	Cl	OH
	Ial.158	NCH ₃	H	H	H	H	Br	OH
25	Ial.159	NCH ₃	H	H	H	H	NO ₂	OH
	Ial.160	NCH ₃	H	H	H	H	CHF ₂	OH
	Ial.161	NCH ₃	H	H	H	H	CF ₃	OH
	Ial.162	NCH ₃	H	H	H	H	OCH ₃	OH
	Ial.163	NCH ₃	H	H	H	H	OCH ₂ CH ₃	OH
30	Ial.164	NCH ₃	H	H	H	H	OCHF ₂	OH
	Ial.165	NCH ₃	H	H	H	H	OCF ₃	OH
	Ial.166	NCH ₃	CH ₃	H	H	H	SCH ₃	OH
	Ial.167	NCH ₃	CH ₃	H	H	H	SCH ₂ CH ₃	OH
	Ial.168	NCH ₃	CH ₃	H	H	H	SO ₂ CH ₃	OH
35	Ial.169	NCH ₃	CH ₃	H	H	H	SO ₂ CH ₂ CH ₃	OH
	Ial.170	NCH ₃	CH ₃	H	H	H	SO ₂ CH(CH ₃) ₂	OH
	Ial.171	NCH ₃	CH ₃	H	H	H	SO ₂ (CH ₂) ₂ CH ₃	OH
	Ial.172	NCH ₃	CH ₃	H	H	H	Cl	OH
	Ial.173	NCH ₃	CH ₃	H	H	H	Br	OH
40	Ial.174	NCH ₃	CH ₃	H	H	H	NO ₂	OH
	Ial.175	NCH ₃	CH ₃	H	H	H	CHF ₂	OH
	Ial.176	NCH ₃	CH ₃	H	H	H	CF ₃	OH
	Ial.177	NCH ₃	CH ₃	H	H	H	OCH ₃	OH
	Ial.177	NCH ₃	CH ₃	H	H	H	OCH ₃	OH

	No.	X	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷
5	Ia1.178	NCH ₃	CH ₃	H	H	H	OCH ₂ CH ₃	OH
	Ia1.179	NCH ₃	CH ₃	H	H	H	OCHF ₂	OH
	Ia1.180	NCH ₃	CH ₃	H	H	H	OCF ₃	OH
	Ia1.181	NCH ₃	H	H	CH ₃	H	SCH ₃	OH
	Ia1.182	NCH ₃	H	H	CH ₃	H	SCH ₂ CH ₃	OH
10	Ia1.183	NCH ₃	H	H	CH ₃	H	SO ₂ CH ₃	OH
	Ia1.184	NCH ₃	H	H	CH ₃	H	SO ₂ CH ₂ CH ₃	OH
	Ia1.185	NCH ₃	H	H	CH ₃	H	SO ₂ CH(CH ₃) ₂	OH
	Ia1.186	NCH ₃	H	H	CH ₃	H	SO ₂ (CH ₂) ₂ CH ₃	OH
	Ia1.187	NCH ₃	H	H	CH ₃	H	Cl	OH
15	Ia1.188	NCH ₃	H	H	CH ₃	H	Br	OH
	Ia1.189	NCH ₃	H	H	CH ₃	H	NO ₂	OH
	Ia1.190	NCH ₃	H	H	CH ₃	H	CHF ₂	OH
	Ia1.191	NCH ₃	H	H	CH ₃	H	CF ₃	OH
	Ia1.192	NCH ₃	H	H	CH ₃	H	OCH ₃	OH
20	Ia1.193	NCH ₃	H	H	CH ₃	H	OCH ₂ CH ₃	OH
	Ia1.194	NCH ₃	H	H	CH ₃	H	OCHF ₂	OH
	Ia1.195	NCH ₃	H	H	CH ₃	H	OCF ₃	OH
	Ia1.196	NCH ₃	CH ₃	CH ₃	H	H	SCH ₃	OH
	Ia1.197	NCH ₃	CH ₃	CH ₃	H	H	SCH ₂ CH ₃	OH
25	Ia1.198	NCH ₃	CH ₃	CH ₃	H	H	SO ₂ CH ₃	OH
	Ia1.199	NCH ₃	CH ₃	CH ₃	H	H	SO ₂ CH ₂ CH ₃	OH
	Ia1.200	NCH ₃	CH ₃	CH ₃	H	H	SO ₂ CH(CH ₃) ₂	OH
	Ia1.201	NCH ₃	CH ₃	CH ₃	H	H	SO ₂ (CH ₂) ₂ CH ₃	OH
	Ia1.202	NCH ₃	CH ₃	CH ₃	H	H	Cl	OH
30	Ia1.203	NCH ₃	CH ₃	CH ₃	H	H	Br	OH
	Ia1.204	NCH ₃	CH ₃	CH ₃	H	H	NO ₂	OH
	Ia1.205	NCH ₃	CH ₃	CH ₃	H	H	CHF ₂	OH
	Ia1.206	NCH ₃	CH ₃	CH ₃	H	H	CF ₃	OH
	Ia1.207	NCH ₃	CH ₃	CH ₃	H	H	OCH ₃	OH
35	Ia1.208	NCH ₃	CH ₃	CH ₃	H	H	OCH ₂ CH ₃	OH
	Ia1.209	NCH ₃	CH ₃	CH ₃	H	H	OCHF ₂	OH
	Ia1.210	NCH ₃	CH ₃	CH ₃	H	H	OCF ₃	OH
	Ia1.211	NCH ₃	CH ₃	H	CH ₃	H	SCH ₃	OH
	Ia1.212	NCH ₃	CH ₃	H	CH ₃	H	SCH ₂ CH ₃	OH
40	Ia1.213	NCH ₃	CH ₃	H	CH ₃	H	SO ₂ CH ₃	OH
	Ia1.214	NCH ₃	CH ₃	H	CH ₃	H	SO ₂ CH ₂ CH ₃	OH
	Ia1.215	NCH ₃	CH ₃	H	CH ₃	H	SO ₂ CH(CH ₃) ₂	OH
	Ia1.216	NCH ₃	CH ₃	H	CH ₃	H	SO ₂ (CH ₂) ₂ CH ₃	OH


	No.	X	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷
5	Ia1.217	NCH ₃	CH ₃	H	CH ₃	H	Cl	OH
	Ia1.218	NCH ₃	CH ₃	H	CH ₃	H	Br	OH
	Ia1.219	NCH ₃	CH ₃	H	CH ₃	H	NO ₂	OH
	Ia1.220	NCH ₃	CH ₃	H	CH ₃	H	CHF ₂	OH
	Ia1.221	NCH ₃	CH ₃	H	CH ₃	H	CF ₃	OH
10	Ia1.222	NCH ₃	CH ₃	H	CH ₃	H	OCH ₃	OH
	Ia1.223	NCH ₃	CH ₃	H	CH ₃	H	OCH ₂ CH ₃	OH
	Ia1.224	NCH ₃	CH ₃	H	CH ₃	H	OCHF ₂	OH
	Ia1.225	NCH ₃	CH ₃	H	CH ₃	H	OCF ₃	OH
	Ia1.226	NCH ₃	H	H	CH ₃	CH ₃	SCH ₃	OH
15	Ia1.227	NCH ₃	H	H	CH ₃	CH ₃	SCH ₂ CH ₃	OH
	Ia1.228	NCH ₃	H	H	CH ₃	CH ₃	SO ₂ CH ₃	OH
	Ia1.229	NCH ₃	H	H	CH ₃	CH ₃	SO ₂ CH ₂ CH ₃	OH
	Ia1.230	NCH ₃	H	H	CH ₃	CH ₃	SO ₂ CH(CH ₃) ₂	OH
	Ia1.231	NCH ₃	H	H	CH ₃	CH ₃	SO ₂ (CH ₂) ₂ CH ₃	OH
20	Ia1.232	NCH ₃	H	H	CH ₃	CH ₃	Cl	OH
	Ia1.233	NCH ₃	H	H	CH ₃	CH ₃	Br	OH
	Ia1.234	NCH ₃	H	H	CH ₃	CH ₃	NO ₂	OH
	Ia1.235	NCH ₃	H	H	CH ₃	CH ₃	CHF ₂	OH
	Ia1.236	NCH ₃	H	H	CH ₃	CH ₃	CF ₃	OH
25	Ia1.237	NCH ₃	H	H	CH ₃	CH ₃	OCH ₃	OH
	Ia1.238	NCH ₃	H	H	CH ₃	CH ₃	OCH ₂ CH ₃	OH
	Ia1.239	NCH ₃	H	H	CH ₃	CH ₃	OCHF ₂	OH
	Ia1.240	NCH ₃	H	H	CH ₃	CH ₃	OCF ₃	OH
	Ia1.241	NCH ₃	CH ₃	CH ₃	CH ₃	H	SCH ₃	OH
30	Ia1.242	NCH ₃	CH ₃	CH ₃	CH ₃	H	SCH ₂ CH ₃	OH
	Ia1.243	NCH ₃	CH ₃	CH ₃	CH ₃	H	SO ₂ CH ₃	OH
	Ia1.244	NCH ₃	CH ₃	CH ₃	CH ₃	H	SO ₂ CH ₂ CH ₃	OH
	Ia1.245	NCH ₃	CH ₃	CH ₃	CH ₃	H	SO ₂ CH(CH ₃) ₂	OH
	Ia1.246	NCH ₃	CH ₃	CH ₃	CH ₃	H	SO ₂ (CH ₂) ₂ CH ₃	OH
35	Ia1.247	NCH ₃	CH ₃	CH ₃	CH ₃	H	Cl	OH
	Ia1.248	NCH ₃	CH ₃	CH ₃	CH ₃	H	Br	OH
	Ia1.249	NCH ₃	CH ₃	CH ₃	CH ₃	H	NO ₂	OH
	Ia1.250	NCH ₃	CH ₃	CH ₃	CH ₃	H	CHF ₂	OH
	Ia1.251	NCH ₃	CH ₃	CH ₃	CH ₃	H	CF ₃	OH
40	Ia1.252	NCH ₃	CH ₃	CH ₃	CH ₃	H	OCH ₃	OH
	Ia1.253	NCH ₃	CH ₃	CH ₃	CH ₃	H	OCH ₂ CH ₃	OH
	Ia1.254	NCH ₃	CH ₃	CH ₃	CH ₃	H	OCHF ₂	OH
	Ia1.255	NCH ₃	CH ₃	CH ₃	CH ₃	H	OCF ₃	OH
	Ia1.255	NCH ₃	CH ₃	CH ₃	CH ₃	H	OCF ₃	OH

	No.	X	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷
5	Ia1.256	NCH ₃	CH ₃	H	CH ₃	CH ₃	SCH ₃	OH
	Ia1.257	NCH ₃	CH ₃	H	CH ₃	CH ₃	SCH ₂ CH ₃	OH
	Ia1.258	NCH ₃	CH ₃	H	CH ₃	CH ₃	SO ₂ CH ₃	OH
	Ia1.259	NCH ₃	CH ₃	H	CH ₃	CH ₃	SO ₂ CH ₂ CH ₃	OH
	Ia1.260	NCH ₃	CH ₃	H	CH ₃	CH ₃	SO ₂ CH(CH ₃) ₂	OH
10	Ia1.261	NCH ₃	CH ₃	H	CH ₃	CH ₃	SO ₂ (CH ₂) ₂ CH ₃	OH
	Ia1.262	NCH ₃	CH ₃	H	CH ₃	CH ₃	Cl	OH
	Ia1.263	NCH ₃	CH ₃	H	CH ₃	CH ₃	Br	OH
	Ia1.264	NCH ₃	CH ₃	H	CH ₃	CH ₃	NO ₂	OH
	Ia1.265	NCH ₃	CH ₃	H	CH ₃	CH ₃	CHF ₂	OH
15	Ia1.266	NCH ₃	CH ₃	H	CH ₃	CH ₃	CF ₃	OH
	Ia1.267	NCH ₃	CH ₃	H	CH ₃	CH ₃	OCH ₃	OH
	Ia1.268	NCH ₃	CH ₃	H	CH ₃	CH ₃	OCH ₂ CH ₃	OH
	Ia1.269	NCH ₃	CH ₃	H	CH ₃	CH ₃	OCHF ₂	OH
	Ia1.270	NCH ₃	CH ₃	H	CH ₃	CH ₃	OCF ₃	OH
20	Ia1.271	NCH ₃	CH ₃	CH ₃	CH ₃	CH ₃	SCH ₃	OH
	Ia1.272	NCH ₃	CH ₃	CH ₃	CH ₃	CH ₃	SCH ₂ CH ₃	OH
	Ia1.273	NCH ₃	CH ₃	CH ₃	CH ₃	CH ₃	SO ₂ CH ₃	OH
	Ia1.274	NCH ₃	CH ₃	CH ₃	CH ₃	CH ₃	SO ₂ CH ₂ CH ₃	OH
	Ia1.275	NCH ₃	CH ₃	CH ₃	CH ₃	CH ₃	SO ₂ CH(CH ₃) ₂	OH
25	Ia1.276	NCH ₃	CH ₃	CH ₃	CH ₃	CH ₃	SO ₂ (CH ₂) ₂ CH ₃	OH
	Ia1.277	NCH ₃	CH ₃	CH ₃	CH ₃	CH ₃	Cl	OH
	Ia1.278	NCH ₃	CH ₃	CH ₃	CH ₃	CH ₃	Br	OH
	Ia1.279	NCH ₃	CH ₃	CH ₃	CH ₃	CH ₃	NO ₂	OH
	Ia1.280	NCH ₃	CH ₃	CH ₃	CH ₃	CH ₃	CHF ₂	OH
30	Ia1.281	NCH ₃	CH ₃	CH ₃	CH ₃	CH ₃	CF ₃	OH
	Ia1.282	NCH ₃	CH ₃	CH ₃	CH ₃	CH ₃	OCH ₃	OH
	Ia1.283	NCH ₃	CH ₃	CH ₃	CH ₃	CH ₃	OCH ₂ CH ₃	OH
	Ia1.284	NCH ₃	CH ₃	CH ₃	CH ₃	CH ₃	OCHF ₂	OH
	Ia1.285	NCH ₃	CH ₃	CH ₃	CH ₃	CH ₃	OCF ₃	OH
35	Ia1.286	NCH ₃	CH ₂ Cl	H	H	H	SCH ₃	OH
	Ia1.287	NCH ₃	CH ₂ Cl	H	H	H	SCH ₂ CH ₃	OH
	Ia1.288	NCH ₃	CH ₂ Cl	H	H	H	SO ₂ CH ₃	OH
	Ia1.289	NCH ₃	CH ₂ Cl	H	H	H	SO ₂ CH ₂ CH ₃	OH
	Ia1.290	NCH ₃	CH ₂ Cl	H	H	H	SO ₂ CH(CH ₃) ₂	OH
40	Ia1.291	NCH ₃	CH ₂ Cl	H	H	H	SO ₂ (CH ₂) ₂ CH ₃	OH
	Ia1.292	NCH ₃	CH ₂ Cl	H	H	H	Cl	OH
	Ia1.293	NCH ₃	CH ₂ Cl	H	H	H	Br	OH
	Ia1.294	NCH ₃	CH ₂ Cl	H	H	H	NO ₂	OH

No.	X	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷
Ia1.295	NCH ₃	CH ₂ Cl	H	H	H	CHF ₂	OH
Ia1.296	NCH ₃	CH ₂ Cl	H	H	H	CF ₃	OH
Ia1.297	NCH ₃	CH ₂ Cl	H	H	H	OCH ₃	OH
Ia1.298	NCH ₃	CH ₂ Cl	H	H	H	OCH ₂ CH ₃	OH
Ia1.299	NCH ₃	CH ₂ Cl	H	H	H	OCHF ₂	OH
Ia1.300	NCH ₃	CH ₂ Cl	H	H	H	OCF ₃	OH

- 10 Extraordinary preference is also given to the compounds of the formula Ia2, in particular to the compounds Ia2.1 to Ia2.300, which differ from the corresponding compounds Ia1.1 to Ia1.300 in that R¹¹ is methyl.

15




20

- Extraordinary preference is also given to the compounds of the formula Ia3, in particular to the compounds Ia3.1 to Ia3.300, which differ from the compounds Ia1.1 to Ia1.300 in that R⁸ is ethyl.

25


30



- Extraordinary preference is also given to the compounds of the formula Ia4, in particular to the compounds Ia4.1 to Ia4.300, which differ from the compounds Ia1.1 to Ia1.300 in that R⁸ is ethyl and R¹¹ is methyl.

35

40




45

25

Extraordinary preference is also given to the compounds of the formula Ia5, in particular to the compounds Ia5.1 to Ia5.300, which differ from the compounds Ia1.1 to Ia1.300 in that R⁸ is 1-methyl-1-ethyl.

5


10



Extraordinary preference is also given to the compounds of the formula Ia6, in particular to the compounds Ia6.1 to Ia6.300, which differ from the compounds Ia1.1 to Ia1.300 in that R⁸ is 1-methyl-1-ethyl and R¹¹ is methyl.


20

25



Extraordinary preference is also given to the compounds of the formula Ia7, in particular to the compounds Ia7.1 to Ia7.300, which differ from the compounds Ia1.1 to Ia1.300 in that R¹⁰ is methyl.

35




Extraordinary preference is also given to the compounds of the formula Ia8, in particular to the compounds Ia8.1 to Ia8.300, which differ from the compounds Ia1.1 to Ia1.300 in that R¹⁰ is methyl and R¹¹ is methyl.

45

26


5



Ia8

Extraordinary preference is also given to the compounds of the
 10 formula Ia9, in particular to the compounds Ia9.1 to Ia9.300,
 which differ from the compounds Ia1.1 to Ia1.300 in that R⁸ and R⁹
 are 1-methyl-1-ethyl.

15




Ia9

20

Extraordinary preference is also given to the compounds of the
 formula Ia10, in particular to the compounds Ia10.1 to Ia10.300,
 which differ from the compounds Ia1.1 to Ia1.300 in that R⁸ and R⁹
 25 are 1-methyl-1-ethyl and R¹¹ is methyl.


30



Ia10

35 Extraordinary preference is also given to the compounds of the
 formula Ia11, in particular to the compounds Ia11.1 to Ia11.300,
 which differ from the compounds Ia1.1 to Ia1.300 in that R⁸ is
 ethyl and R¹⁰ is methyl.

40



Ia11

45

5
10
15
20
25
30
35
40
45

10



Process A:


25

30

35

40

45



L¹ is a nucleophilically replaceable leaving group, such as halogen, for example bromine, chlorine, hetaryl, for example imidazolyl, pyridyl, carboxylate, for example acetate,
5 trifluoroacetate etc.

The activated benzoic acid can be employed directly, such as in the case of the benzoyl halides, or be generated in situ, for example using dicyclohexylcarbodiimide,
10 triphenylphosphine/azodicarboxylic ester, 2-pyridine disulfide/triphenylphosphine, carbonyldiimidazole, etc.

It may be advantageous to carry out the acylation reaction in the presence of a base. The reactants and the auxiliary base are
15 advantageously employed in equimolar amounts. A slight excess of auxiliary base, for example from 1.2 to 1.5 molar equivalents, based on II, may be advantageous in certain cases.

Suitable auxiliary bases are tertiary alkylamines, pyridine or
20 alkali metal carbonates. Suitable for use as solvents are, for example, chlorinated hydrocarbons, such as methylene chloride, 1,2-dichloroethane, aromatic hydrocarbons, such as toluene, xylene, chlorobenzene, ethers, such as diethyl ether, methyl tert-butyl ether, dimethoxyethane, tetrahydrofuran, dioxane,
25 polar aprotic solvents, such as acetonitrile, dimethylformamide, dimethyl sulfoxide, or esters, such as ethyl acetate, or mixtures of these.

If the activated carboxylic acid component used is a benzoyl
30 halide, it may be advantageous to cool the reaction mixture to 0 - 10°C when adding this reaction partner. The mixture is subsequently stirred at 20 - 100°C, preferably at 25 - 50°C, until the reaction has ended. Work-up is carried out in a customary manner, for example by pouring the reaction mixture into water
35 and extracting the product of value. Solvents which are particularly suitable for this purpose are methylene chloride, diethyl ether, dimethoxyethane and ethyl acetate. The organic phase is dried and the solvent is removed, after which the crude ester can be employed for the rearrangement without any further
40 purification.

The rearrangement of the esters to the compounds of the formula I is advantageously carried out at 20 - 40°C in a solvent and in the presence of a base and, if appropriate, using a cyano compound as
45 catalyst.

30

Suitable solvents are, for example, acetonitrile, methylene chloride, 1,2-dichloroethane, dioxane, ethyl acetate, dimethoxyethane, toluene or mixtures of these. Preferred solvents are acetonitrile and dioxane.

5

Suitable bases are tertiary amines, such as triethylamine or pyridine, or alkali metal carbonates, such as sodium carbonate or potassium carbonate, which are preferably employed in an equimolar amount or an up to four-fold excess, based on the ester. Preference is given to using triethylamine or alkali metal carbonates, preferably in twice the equimolar amount, based on the ester.

Suitable cyano compounds are inorganic cyanides, such as sodium cyanide and potassium cyanide, and organic cyano compounds, such as acetonecyanohydrine and trimethylsilyl cyanide. They are employed in an amount of from 1 to 50 mol percent, based on the ester. Preference is given to using acetonecyanohydrine or trimethylsilyl cyanide, for example in an amount of from 5 to 15, preferably 10, mol percent, based on the ester.


Work-up can be carried out in the manner known per se. The reaction mixture is, for example, acidified with dilute mineral acid, such as 5% strength hydrochloric acid or sulfuric acid, and extracted with an organic solvent, for example methylene chloride or ethyl acetate. The organic extract can be extracted with 5-10% strength alkali metal carbonate solution, for example sodium carbonate or potassium carbonate solution. The aqueous phase is acidified and the resulting precipitate is filtered off with suction and/or extracted with methylene chloride or ethyl acetate, and the mixture is dried and concentrated.

(Examples for the preparation of esters of hydroxypyrazoles and for the rearrangement of the esters are given, for example, in EP-A 282 944 and US 4 643 757).

35

However, it is also possible to generate the "acylation product" IV in situ by reacting a pyrazole of the formula II, or an alkali metal salt thereof, with a 3-(heterocyclyl)benzene derivative of the formula V in the presence of carbon monoxide, a catalyst and a base.

45



L^2 is a leaving group, such as halogen, for example chlorine, bromine or iodine, or sulfonate, such as mesylate or triflate; preference is given to bromine or triflate.

The "acylation product" IV proceeds to react, directly or indirectly, to give the 3-(heterocyclyl)-substituted benzoylpyrazole of the formula I.

Suitable catalysts are palladium-ligand complexes in which the palladium is present in oxidation state 0, metallic palladium, which has optionally been absorbed on a carrier, and preferably palladium(II) salts. The reaction with palladium(II) salts and metallic palladium is preferably carried out in the presence of complex ligands.

An example of a suitable palladium(0)-ligand complex is tetrakis(triphenylphosphine)palladium.



Metallic palladium is preferably absorbed on an inert carrier such as, for example, activated carbon, silica, alumina, barium sulfate or calcium carbonate. The reaction is preferably carried out in the presence of complex ligands such as, for example, triphenylphosphine.

32

Examples of suitable palladium(II) salts are palladium acetate and palladium chloride. The presence of complex ligands such as, for example, triphenylphosphine is preferred.

- 5 Suitable complex ligands for the palladium-ligand complexes, or in whose presence the reaction is preferably carried out with metallic palladium or palladium(II) salts, are tertiary phosphines whose structure is represented by the following formulae:

10



- 15 where z is 1 to 4 and the radicals R^{a} to R^{g} are C_1 - C_6 -alkyl, C_3 - C_6 -cycloalkyl, aryl- C_1 - C_2 -alkyl or, preferably, aryl. Aryl is, for example, naphthyl and unsubstituted or substituted phenyl such as, for example, 2-tolyl and, in particular, unsubstituted phenyl.

20

The complex palladium salts can be prepared in a manner known per se starting from commercially available palladium salts such as palladium chloride or palladium acetate and the appropriate phosphines such as, for example, triphenylphosphine or

- 25 1,2-bis(diphenylphosphino)ethane. Many of the complexed palladium salts are also commercially available. Preferred palladium salts are [(R)(+)-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl]-palladium(II) chloride, bis(triphenylphosphine)palladium(II) acetate and, in particular, bis(triphenylphosphine)palladium(II) chloride.

The palladium catalyst is usually employed in a concentration of from 0.05 to 5 mol%, and preferably 1-3 mol%.

- 35 Suitable bases are tertiary amines, such as, for example, N-methylpiperidine, ethyldiisopropylamine, 1,8-bisdimethylaminonaphthalene or, in particular, triethylamine. Also suitable are alkali metal carbonates, such as sodium carbonate or potassium carbonate. However, mixtures of potassium
- 40 carbonate and triethylamine are also suitable.

In general, from 2 to 4 molar equivalent, in particular 2 molar equivalents, of the alkali metal carbonate, and from 1 to 4 molar equivalents, in particular 2 molar equivalents, of the tertiary


45 amine are employed, based on the 3-(heterocyclyl)-benzene derivatives of the formula V.

33

Suitable solvents are nitriles, such as benzonitrile and acetonitrile, amides, such as dimethylformamide, dimethylacetamide, tetra-C₁-C₄-alkylureas or N-methylpyrrolidone and, preferably, ethers, such as tetrahydrofuran and methyl tert-butyl ethers. Particular preference is given to ethers, such as 1,4-dioxane and dimethoxyethane.

Process B:

- 10 Compounds of the formula I where R⁷ ≠ hydroxyl are obtained by reacting compounds of the formula I where R⁷ = hydroxyl with alkylating agents, sulfonylating agents or acylating agents L³-R^{7a} (VI).



- 25 L³ is a nucleophilically replaceable leaving group, such as halogen, for example bromine or chlorine, acyloxy, for example acetyloxy or ethylcarbonyloxy, or alkylsulfonyloxy, for example methylsulfonyloxy or trifluoromethylsulfonyloxy.
- 30 R^{7a} is C₁-C₆-alkyl, C₃-C₆-alkenyl, C₁-C₆-alkylsulfonyl, C₁-C₆-alkylcarbonyl, C₁-C₄-(alkylthio)carbonyl, phenylsulfonyl or phenylcarbonyl, where the phenyl radical of the two last-mentioned substituents may be partially or fully halogenated and/or may carry one to three of the following groups:
- 35 nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy or C₁-C₄-haloalkoxy.

The compounds of the formula VI can be employed directly, such as, for example, in the case of the sulfonyl halides or sulfonic anhydrides, or be generated in situ, for example activated sulfonic acids (using sulfonic acid and dicyclohexylcarbonyldiimide, carbonyldiimidazole, etc.).

The starting materials are generally employed in equimolar amounts. However, it may also be advantageous to employ an excess of one or the other component.

- 5 If appropriate, it may be advantageous to carry out the reaction in the presence of a base. The reactants and the auxiliary base are advantageously employed in equimolar amounts. An excess of auxiliary base, for example from 1.5 to 3 molar equivalents, based on I, may be advantageous in certain cases.

- 10 Suitable auxiliary bases are tertiary alkylamines, such as triethylamine, pyridine, alkali metal carbonates, for example sodium carbonate or potassium carbonate, and alkali metal hydrides, for example sodium hydride. Preference is given to using triethylamine and pyridine.


- Suitable solvents are, for example, chlorinated hydrocarbons, such as methylene chloride and 1,2-dichloroethane, aromatic hydrocarbons, for example toluene, xylene, chlorobenzene, ethers, 20 such as diethyl ether, methyl tert-butyl ether, tetrahydrofuran and dioxane, polar aprotic solvents, such as acetonitrile, dimethylformamide, dimethyl sulfoxide, or esters, such as ethyl acetate, or mixtures of these.

- 25 In general, the reaction temperature is in the range from 0°C to the boiling point of the reaction mixture.

Work-up can be carried out in the manner known per se to give the product.

- 30 The pyrazoles of the formula II used as starting materials are known or can be prepared by the process known per se (for example EP-A 240 001 and J. Prakt. Chem. 315, 383 (1973)).

- 35 The activated benzoic acids III α can be obtained in a manner known per se from the benzoic acids III β . The latter for their part are obtained by hydrolysis from the corresponding esters VII. These can be prepared by converting an oxime or hydrazone of the formula VIII into the corresponding hydroxamic acid halide, 40 in particular hydroxamic acid chloride, or carbohydrazide halide, in particular carbohydrazide chloride; generating a nitrile oxide or nitrile imine in situ and reacting this with an alkene (cf., for example, Chem. Ber. 106, 3258-3274 (1973)).



L⁴ denotes a C₁-C₆-alkoxy radical.

25 However, the benzoic acids III β can also be obtained by converting an oxime or hydrazine of the formula IX into the corresponding nitrile oxides or nitrile imines and reacting these with alkenes to give the corresponding cycloaddition products (cf., for example, Chem. Ber. 106, 3258-3274 (1973)). Thus, for example, the


30 oxime of the formula IX (X=O) is oxidized with sodium hypochlorite and reacted with a suitable alkene in an inert solvent such as methylene chloride, chloroform, tetrahydrofuran, dioxane or acetonitrile. The product is then converted in the presence of a catalyst and a base into the benzoic acid III β using carbon monoxide

35 and water.

40

45


36



L² denotes a leaving group, such as halogen, for example chlorine, bromine or iodine, or sulfonate, such as mesylate or triflate, preferably bromine or triflate.

With respect to the carbonylation reaction, what has been said above applies analogously.

The compounds of the formulae III and V are in each case novel as such



where in each case the variables R¹, R³ to R⁶ and X are as defined under the compounds of the formula I and

45

R² is C₁-C₄-haloalkyl;
 L is hydroxyl or a radical that can be removed by hydrolysis;
 or
 L² is a leaving group that can be displaced nucleophilically.

5

Examples of radicals that can be removed by hydrolysis are alkoxy, phenoxy, alkylthio and phenylthio radicals, which may be unsubstituted or substituted, halides, hetaryl radicals attached via nitrogen, amino and imino radicals, which may be unsubsti-

10

tuted or substituted, etc.

Examples of nucleophilically displaceable leaving groups are halogen, C₁-C₄-alkylsulfonyloxy and C₁-C₄-haloalkylsulfonyloxy;

15 Preferred compounds of the formula III are those compounds in which L is halogen, in particular chlorine or bromine.

Preference is also given to those compounds of the formula III in which L is C₁-C₆-alkoxy.

20

Preference is also given to those compounds of the formula III in which L is hydroxyl.

With respect to the variables X, R¹ and R³ to R⁶, the particularly
 25 preferred embodiments of the compounds of the formulae III and V correspond to those of the compounds of the formula I.

Preparation examples

30 4-[2-Methyl-3-(4,5-dihydroisoxazol-3-yl)-4-methylsulfonylbenzoyl]-5-hydroxy-1-(1,1-dimethyl-1-ethyl)-1H-pyrazole
 (Compound 2.1)

Variant 1:

35

0.61 g (3.5 mmol) of 1-(1,1-dimethyl-1-ethyl)-5-hydroxy-1H-pyrazole and 0.79 g (3.9 mmol) of dicyclohexylcarbodiimide were added to a solution of 1.0 g (3.5 mmol) of 2-methyl-3-(4,5-dihydroisoxazol-3-yl)-4-methylsulfonylbenzoic acid in
 40 dioxane, and the mixture was stirred at room temperature for 12 hours. Insoluble components were separated off, and 0.58 g (4.2 mmol) of potassium carbonate was then added and the mixture was refluxed for 3 hours. The solvent was then distilled off, the residue was taken up in 5% strength potassium carbonate solution
 45 and washed with methylene chloride and toluene, the pH was adjusted to 3 using 10% strength hydrochloric acid and the mixture was extracted with methylene chloride. This methylene

38

chloride solution was dried and the solvent was removed. This gave 0.81 g (57% of theory) of 4-[2-methyl-3-(4,5-dihydroisoxazol-3-yl)-4-methylsulfonylbenzoyl]-5-hydroxy-1-(1,1-dimethyl-1-ethyl)-1H-pyrazole]

5 M.p. 195-198°C.

Variant 2

- 3.78 g (32 mmol) of thionyl chloride were added to a solution of
10 6 g (21.2 mmol) of 2-methyl-3-(4,5-dihydroisoxazol-3-yl)-4-methylsulfonylbenzoic acid in toluene, and the mixture was refluxed for 4 hours. The solvent was removed, the residue was then taken up in dimethoxyethane and a solution of 2.97 g (21.2 mmol) of 1-(1,1-dimethyl-1-ethyl)-5-hydroxy-1H-pyrazole in
15 dimethoxyethane was added dropwise. The mixture was subsequently stirred at room temperature for 12 hours and refluxed for 3.5 hours. The reaction mixture was cooled, the solvent was distilled off, and the residue was taken up in 5% strength potassium carbonate solution and washed with methylene chloride. The
20 resulting aqueous phase was adjusted to pH = 1 using hydrochloric acid, and the resulting precipitate was filtered off with suction. This gave 4.89 g (52%) of 4-[2-methyl-3-(4,5-dihydroisoxazol-3-yl)-4-methylsulfonylbenzoyl]-5-hydroxy-1-(1,1-dimethyl-1-ethyl)-1H-pyrazole]
25 M.p. 195-198°C.

In addition to the compound above, Table 2 lists further 3-(heterocyclyl)-substituted benzoylpyrazoles of the formula I which were prepared or are preparable in a similar manner.

30

35

40

45




Table 2:

No.	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	R ⁸	R ⁹	R ¹⁰	R ¹¹	physical data m.p. [°C] ¹ H-NMR [δ in ppm]
2.1	CH ₃	H	H	H	H	SO ₂ CH ₃	OH	CH ₃	CH ₃	CH ₃	H	195-198
2.2	CH ₃	H	H	H	H	SO ₂ CH ₃	OCH(CH ₃) ₂	CH ₃	CH ₃	CH ₃	H	oil
2.3	CH ₃	H	H	H	H	SO ₂ CH ₃	OCO[3-F-C ₆ H ₄]	CH ₃	CH ₃	CH ₃	H	210-211
2.4	CH ₃	H	H	H	H	SO ₂ CH ₃	OCO[3,5-(CF ₃) ₂ -C ₆ H ₃]	CH ₃	CH ₃	CH ₃	H	oil
2.5	CH ₃	H	H	H	H	SO ₂ CH ₃	OCO[2,4-F ₂ -C ₆ H ₃]	CH ₃	CH ₃	CH ₃	H	145-148
2.6	CH ₃	H	H	H	H	SO ₂ CH ₃	OCO[3,5-F ₂ -C ₆ H ₃]	CH ₃	CH ₃	CH ₃	H	100-105
2.7	CH ₃	H	H	H	H	SO ₂ CH ₃	OCO[3-(CF ₃)-C ₆ H ₄]	CH ₃	CH ₃	CH ₃	H	90-93
2.8	CH ₃	H	H	H	H	SO ₂ CH ₃	OH	CH ₃	CH ₃	CH ₃	CH ₃	198-200
2.9	CH ₃	H	H	H	H	SO ₂ CH ₃	OH	CH ₃	CH ₃	H	H	203-205
2.10	CH ₂ -CH ₃	H	H	H	H	SO ₂ CH ₃	OH	CH ₃	CH ₃	CH ₃	H	75-80

No.	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	R ⁸	R ⁹	R ¹⁰	R ¹¹	physical data m.p. [°C] ¹ H-NMR [δ in ppm]
2.11	CH ₃	H	H	H	H	SO ₂ CH ₃	OH	CH(CH ₃) ₂	CH ₃	H	H	90-95
2.12	CH ₃	H	H	H	H	SO ₂ CH ₃	OCOC ₆ H ₅	CH(CH ₃) ₂	CH ₃	H	H	95-100
2.13	CH ₃	CH ₂ Cl	H	H	H	SO ₂ CH ₃	OH	CH ₃	CH ₃	CH ₃	H	80-81
2.14	CH ₃	CH ₂ Cl	H	H	H	SO ₂ CH ₃	OH	CH ₃	CH ₃	H	H	75-84
2.15	CH ₃	CH ₂ Cl	H	H	H	SO ₂ CH ₃	OCO[3-F-C ₆ H ₄]	CH ₃	CH ₃	H	H	77-83
2.16	CH ₃	H	H	H	H	SO ₂ CH ₃	OH	CH(CH ₃) ₂	CH(CH ₃) ₂	H	H	120-125
2.17	CH ₃	CH ₂ Cl	CH ₂ Cl	H	H	SO ₂ CH ₃	OH	CH ₃	CH ₃	H	H	73-79
2.18	CH ₃	CH ₂ Cl	CH ₂ Cl	H	H	SO ₂ CH ₃	OH	CH ₃	CH ₃	CH ₃	H	73-78
2.19	CH ₃	H	H	H	H	SO ₂ CH ₃	OCH ₃	CH ₃	CH ₃	H	H	oil
2.20	CH ₃	H	H	H	H	SO ₂ CH ₃	OCH ₂ CH ₃	CH ₃	CH ₃	H	H	165-170
2.21	CH ₃	H	H	H	H	SO ₂ CH ₃	OCH(CH ₃) ₂	CH ₃	CH ₃	H	H	oil
2.22	CH ₃	H	H	H	H	SO ₂ CH ₃	OCOC ₆ H ₅	CH ₃	CH ₃	H	H	oil
2.23	CH ₃	H	H	H	H	SO ₂ CH ₃	OCO[3-F-C ₆ H ₄]	CH ₃	CH ₃	H	H	110-115
2.24	CH ₃	H	H	H	H	SO ₂ CH ₃	OCOCH ₃	CH ₃	CH ₃	H	H	80-85
2.25	CH ₃	H	H	H	H	SO ₂ CH ₃	OSO ₂ [4-CH ₃ -C ₆ H ₄]	CH ₃	CH ₃	H	H	95-100
2.26	CH ₃	H	H	H	H	SO ₂ CH ₃	OCOSCH ₃	CH ₃	CH ₃	H	H	143-145
2.27	CH ₃	H	H	H	H	SO ₂ CH ₃	OCOC ₆ H ₅	CH ₃	CH ₃	CH ₃	H	228-230

The 3-(heterocyclyl)-substituted benzoylpyrazoles of the formula I and their agriculturally useful salts are suitable, both in the form of isomer mixtures and in the form of the pure isomers, as
 5 herbicides. The herbicidal compositions comprising compounds of the formula I control vegetation on non-crop areas very efficiently, especially at high rates of application. They act against weeds and harmful grasses in crops such as wheat, rice, maize, soya and cotton without causing any significant damage to
 10 the crop plants. This effect is mainly observed at low rates of application.

Depending on the particular application method used, the compounds of the formula I, or the herbicidal compositions
 15 comprising them, can additionally be employed in a further number of crop plants for eliminating undesirable plants. Examples of suitable crops are the following:

Allium cepa, Ananas comosus, Arachis hypogaea, Asparagus
 20 officinalis, Beta vulgaris spec. altissima, Beta vulgaris spec. rapa, Brassica napus var. napus, Brassica napus var. napobrassica, Brassica rapa var. silvestris, Camellia sinensis, Carthamus tinctorius, Carya illinoensis, Citrus limon, Citrus sinensis, Coffea arabica (Coffea canephora, Coffea liberica),
 25 Cucumis sativus, Cynodon dactylon, Daucus carota, Elaeis guineensis, Fragaria vesca, Glycine max, Gossypium hirsutum, (Gossypium arboreum, Gossypium herbaceum, Gossypium vitifolium), Helianthus annuus, Hevea brasiliensis, Hordeum vulgare, Humulus lupulus, Ipomoea batatas, Juglans regia, Lens culinaris, Linum
 30 usitatissimum, Lycopersicon lycopersicum, Malus spec., Manihot esculenta, Medicago sativa, Musa spec., Nicotiana tabacum (N. rustica), Olea europaea, Oryza sativa, Phaseolus lunatus, Phaseolus vulgaris, Picea abies, Pinus spec., Pisum sativum, Prunus avium, Prunus persica, Pyrus communis, Ribes sylvestre,
 35 Ricinus communis, Saccharum officinarum, Secale cereale, Solanum tuberosum, Sorghum bicolor (s. vulgare), Theobroma cacao, Trifolium pratense, Triticum aestivum, Triticum durum, Vicia faba, Vitis vinifera and Zea mays.

40 In addition, the compounds of the formula I may also be used in crops which tolerate the action of herbicides owing to breeding, including genetic engineering methods.

The compounds of the formula I, or the herbicidal compositions
 45 comprising them, can be used for example in the form of ready-to-spray aqueous solutions, powders, suspensions, also highly-concentrated aqueous, oily or other suspensions or

dispersions, emulsions, oil dispersions, pastes, dusts, materials for broadcasting, or granules, by means of spraying, atomizing, dusting, broadcasting or watering. The use forms depend on the intended aims; in any case, they should guarantee a very fine
5 distribution of the active compounds according to the invention.

The herbicidal compositions comprise a herbicidally effective amount of at least one compound of the formula I or an agriculturally useful salt of I and auxiliaries which are
10 customarily used for formulating crop protection agents.

Essentially, suitable inert auxiliaries include:
mineral oil fractions of medium to high boiling point, such as kerosene and diesel oil, furthermore coal tar oils and oils of
15 vegetable or animal origin, aliphatic, cyclic and aromatic hydrocarbons, eg. paraffins, tetrahydronaphthalene, alkylated naphthalenes and their derivatives, alkylated benzenes and their derivatives, alcohols such as methanol, ethanol, propanol, butanol and cyclohexanol, ketones such as cyclohexanone, or
20 strongly polar solvents, e.g. amines such as N-methylpyrrolidone, and water.

Aqueous use forms can be prepared from emulsion concentrates, suspensions, pastes, wettable powders or water-dispersible
25 granules by adding water. To prepare emulsions, pastes or oil dispersions, the substances, either as such or dissolved in an oil or solvent, can be homogenized in water by means of a wetting agent, tackifier, dispersant or emulsifier. Alternatively, it is possible to prepare concentrates comprising active compound,
30 wetting agent, tackifier, dispersant or emulsifier and, if desired, solvent or oil, which are suitable for dilution with water.

Suitable surfactants (adjuvants) are the alkali metal salts,
35 alkaline earth metal salts and ammonium salts of aromatic sulfonic acids, eg. ligno-, phenol-, naphthalene- and dibutyl-naphthalenesulfonic acid, and of fatty acids, alkyl- and alkylarylsulfonates, alkyl sulfates, lauryl ether sulfates and fatty alcohol sulfates, and salts of sulfated hexa-, hepta- and
40 octadecanols, and also of fatty alcohol glycol ether, condensates of sulfonated naphthalene and its derivatives with formaldehyde, condensates of naphthalene, or of the naphthalenesulfonic acids with phenol and formaldehyde, polyoxyethylene octylphenol ether, ethoxylated isooctyl-, octyl- or nonylphenol, alkylphenyl or
45 tributylphenyl polyglycol ether, alkylaryl polyether alcohols, isotridecyl alcohol, fatty alcohol/ethylene oxide condensates, ethoxylated castor oil, polyoxyethylene alkyl ethers or

polyoxypropylene alkyl ethers, lauryl alcohol polyglycol ether acetate, sorbitol esters, lignin-sulfite waste liquors or methylcellulose.

- 5 Powders, materials for broadcasting and dusts can be prepared by mixing or grinding the active compounds together with a solid carrier.

- Granules, e.g. coated granules, impregnated granules and
- 10 homogeneous granules, can be prepared by binding the active compounds to solid carriers. Solid carriers are mineral earths, such as silicas, silica gels, silicates, talc, kaolin, limestone, lime, chalk, bole, loess, clay, dolomite, diatomaceous earth, calcium sulfate, magnesium sulfate, magnesium oxide, ground
- 15 synthetic materials, fertilizers such as ammonium sulfate, ammonium phosphate and ammonium nitrate, ureas, and products of vegetable origin, such as cereal meal, tree bark meal, wood meal and nutshell meal, cellulose powders, or other solid carriers.
- 20 The concentrations of the compounds of the formula I in the ready-to-use preparations can be varied within wide ranges. In general, the formulations comprise about from 0.001 to 98% by weight, preferably 0.01 to 95% by weight, of at least one active compound. The active compounds are employed in a purity of from
- 25 90% to 100%, preferably 95% to 100% (according to the NMR spectrum).

The following formulation examples illustrate the preparation of such formulations:

30

- I. 20 parts by weight of the compound No. 2.8 are dissolved in a mixture composed of 80 parts by weight of alkylated benzene, 10 parts by weight of the adduct of 8 to 10 mol of ethylene oxide to 1 mol of oleic acid N-monoethanolamide, 5
- 35 parts by weight of calcium dodecylbenzenesulfonate and 5 parts by weight of the adduct of 40 mol of ethylene oxide to 1 mol of castor oil. Pouring the solution into 100,000 parts by weight of water and finely distributing it therein gives an aqueous dispersion which comprises 0.02% by weight
- 40 of the active compound.

- II. 20 parts by weight of the compound No. 2.9 are dissolved in a mixture composed of 40 parts by weight of cyclohexanone, 30 parts by weight of isobutanol, 20 parts by weight of
- 45 the adduct of 7 mol of ethylene oxide to 1 mol of isooctylphenol and 10 parts by weight of the adduct of 40 mol of ethylene oxide to 1 mol of castor oil. Pouring

the solution into 100,000 parts by weight of water and finely distributing it therein gives an aqueous dispersion which comprises 0.02% by weight of the active compound.

- 5 III. 20 parts by weight of the active compound No. 2.3 are dissolved in a mixture composed of 25 parts by weight of cyclohexanone, 65 parts by weight of a mineral oil fraction of boiling point 210 to 280°C and 10 parts by weight of the adduct of 40 mol of ethylene oxide to 1 mol of castor oil.
- 10 Pouring the solution into 100,000 parts by weight of water and finely distributing it therein gives an aqueous dispersion which comprises 0.02% by weight of the active compound.
- 15 IV. 20 parts by weight of the active compound No. 2.3 are mixed thoroughly with 3 parts by weight of sodium diisobutyl naphthalenesulfonate, 17 parts by weight of the sodium salt of a lignosulfonic acid from a sulfite waste liquor and 60 parts by weight of pulverulent silica gel,
- 20 and the mixture is ground in a hammer mill. Finely distributing the mixture in 20,000 parts by weight of water gives a spray mixture which comprises 0.1% by weight of the active compound.
- 25 V. 3 parts by weight of the active compound No. 2.9 are mixed with 97 parts by weight of finely divided kaolin. This gives a dust which comprises 3% by weight of the active compound.
- 30 VI. 20 parts by weight of the active compound No. 2.8 are mixed intimately with 2 parts by weight of calcium dodecylbenzenesulfonate, 8 parts by weight of fatty alcohol polyglycol ether, 2 parts by weight of the sodium salt of a phenol/urea/formaldehyde condensate and 68 parts by weight
- 35 of a paraffinic mineral oil. This gives a stable oily dispersion.
- VII. 1 part by weight of the active compound No. 2.3 is dissolved in a mixture composed of 70 parts by weight of cyclohexanone, 20 parts by weight of ethoxylated
- 40 isooctylphenol and 10 parts by weight of ethoxylated castor oil. This gives a stable emulsion concentrate.
- VIII. 1 part by weight of the active compound No. 2.9 is
- 45 dissolved in a mixture composed of 80 parts by weight of cyclohexanone and 20 parts by weight of Wettol® EM 31 (= nonionic emulsifier based on ethoxylated castor oil).

This gives a stable emulsion concentrate.

The compounds of the formula I or the herbicidal compositions can be applied pre- or post-emergence. If the active compounds are less well tolerated by certain crop plants, application techniques may be used in which the herbicidal compositions are sprayed, with the aid of the spraying equipment, in such a way that they come into contact as little as possible, if at all, with the leaves of the sensitive crop plants, while the active compounds reach the leaves of undesirable plants growing underneath, or the bare soil surface (post-directed, lay-by).

The application rates of the compound of the formula I are from 0.001 to 3.0, preferably 0.01 to 1.0, kg/ha of active substance (a.s.), depending on the control target, the season, the target plants and the growth stage.

To widen the activity spectrum and to achieve synergistic effects, the 3-(heterocyclyl)-substituted benzoylpyrazoles of the formula I may be mixed with a large number of representatives of other herbicidal or growth-regulating active compound groups and then applied concomitantly. Suitable components for mixtures are, for example, 1,2,4-thiadiazoles, 1,3,4-thiadiazoles, amides, aminophosphoric acid and its derivatives, aminotriazoles, anilides, (het)aryloxyalkanoic acids and their derivatives, benzoic acid and its derivatives, benzothiadiazinones, 2-(het)aroyl-1,3-cyclohexanediones, hetaryl aryl ketones, benzylisoxazolidinones, meta-CF₃-phenyl derivatives, carbamates, quinolinecarboxylic acid and its derivatives, chloroacetanilides, cyclohexeneone oxime ether derivatives, diazines, dichloropropionic acid and its derivatives, dihydrobenzofurans, dihydrofuran- 3-ones, dinitroanilines, dinitrophenols, diphenyl ethers, dipyridyls, halocarboxylic acids and their derivatives, ureas, 3-phenyluracils, imidazoles, imidazolinones, N-phenyl-3,4,5,6-tetrahydrophthalimides, oxadiazoles, oxiranes, phenols, aryloxy- and hetaryloxyphenoxypropionic esters, phenylacetic acid and its derivatives, 2-phenylpropionic acid and its derivatives, pyrazoles, phenylpyrazoles, pyridazines, pyridinecarboxylic acid and its derivatives, 2-pyrimidyl ethers, sulfonamides, sulfonylureas, triazines, triazinones, triazolinones, triazolecarboxamides and uracils.

It may furthermore be advantageous to apply the compounds of the formula I, alone or else concomitantly in combination with other herbicides, in the form of a mixture with other crop protection agents, for example together with agents for controlling pests or phytopathogenic fungi or bacteria. Also of interest is the

miscibility with mineral salt solutions, which are employed for treating nutritional and trace element deficiencies.

Non-phytotoxic oils and oil concentrates may also be added.

5 Use Examples

The herbicidal activity of the 3-(heterocyclyl)-substituted benzoylpyrazoles of the formula I was demonstrated by the following greenhouse experiments:

10

The culture containers used were plastic flowerpots containing loamy sand with approximately 3.0% of humus as the substrate. The seeds of the test plants were sown separately for each species.

- 15 For the pre-emergence treatment, the active compounds, which had been suspended or emulsified in water, were applied directly after sowing by means of finely distributing nozzles. The containers were irrigated gently to promote germination and growth and subsequently covered with transparent plastic hoods
- 20 until the plants had rooted. This cover causes uniform germination of the test plants, unless this has been adversely effected by the active compounds.

- For the post-emergence treatment, the test plants were first
- 25 grown to a height of 3 to 15 cm, depending on the plant habit, and only then treated with the active compounds which had been suspended or emulsified in water. The test plants were, for this purpose, either sown directly and grown in the same containers, or they were first grown separately as seedlings and transplanted
- 30 into the test containers a few days prior to treatment. The application rate for the post-emergence treatment was 0.5, 0.25, 0.125 or 0.0625 kg of a.s. (active substance)/ha.

- Depending on the species, the plants were kept at 10 - 25°C or
- 35 20 - 35°C . The test period extended over 2 to 4 weeks. During this time, the plants were tended, and their response to the individual treatments was evaluated.

- The evaluation was carried out using a scale from 0 to 100. 100
- 40 means no emergence of the plants, or complete destruction of at least the aerial parts and 0 means no damage, or normal course of growth.

- The plants used in the greenhouse experiments were of the
- 45 following species:

	Scientific name	Common name
	<i>Abutilon theophrasti</i>	velvet leaf
	<i>Amaranthus retroflexus</i>	pigweed
	<i>Avena fatua</i>	wild oat
5	<i>Chenopodium album</i>	lamb's-quarters
	<i>Echinochloa crus galli</i>	barnyard grass
	<i>Polygonum persicaria</i>	lady's-thumb
	<i>Setaria faberi</i>	giant foxtail
	<i>Setaria viridis</i>	green foxtail
	<i>Sinapis alba</i>	white mustard
10	<i>Solanum nigrum</i>	black nightshade

At application rates of 0.125 or 0.0625 kg/ha, the compound 2.3 (Table 2) showed very good action against the undesirable plants barnyard grass, pigweed, lamb's-quarters, lady's-thumb and black
15 nightshade when used in the post-emergence method.

Furthermore, compound 2.8 (Table 2) effected very good post-emergence control at application rates of 0.5 or 0.25 kg/ha of the harmful plants barnyard grass, green foxtail, lamb's-
20 quarters and black nightshade.

The action of compound 2.9 (Table 2), when applied in the post-emergence method at application rates of 0.25 or 0.125 kg/ha, on the grasses wild oat and green foxtail and on the
25 weeds pigweed, lady's-thumb and white mustard is very good.

Likewise, the compound 2.10 (Table 2) effects, at application rates of 0.25 or 0.125 kg/ha, very good post-emergence control of the undesirable plants barnyard grass, lamb's-quarters, white
30 mustard and black nightshade.


35

40

45

We claim:

1. A 3-(heterocyclyl)-substituted benzoylpyrazole of the formula
 5 I



where:

- 15 X is O, NH or N(C₁-C₆-alkyl);
- R¹ is C₁-C₆-alkyl;
- 20 R², R³, R⁴, R⁵ are hydrogen, C₁-C₄-alkyl or C₁-C₄-haloalkyl;
- R⁶ is halogen, nitro, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio, C₁-C₄-alkylsulfonyl or C₁-C₄-haloalkylsulfonyl;
- 25 R⁷ is hydroxyl, C₁-C₆-alkoxy, C₃-C₆-alkenyloxy, C₁-C₆-alkylsulfonyloxy, C₁-C₆-alkylcarbonyloxy, C₁-C₄-(alkylthio)carbonyloxy, phenylsulfonyloxy or phenylcarbonyloxy, where the phenyl radical of the two last-mentioned substituents may be partially or fully halogenated and/or may carry one to three of the following groups:
- 30 nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy or C₁-C₄-haloalkoxy;
- 35 R⁸, R⁹ are C₁-C₄-alkyl;
- R¹⁰ is hydrogen or C₁-C₄-alkyl;
- 40 where the number of the carbon atoms of the radicals R⁸, R⁹ and R¹⁰ together is at most 7,
- R¹¹ is hydrogen or C₁-C₄-alkyl;
- 45 and its agriculturally useful salts.

2. A 3-(heterocyclyl)-substituted benzoylpyrazole of the formula I as claimed in claim 1 where

X is O;

5

R¹ is C₁-C₄-alkyl;

R⁶ is C₁-C₄-alkylthio or C₁-C₄-alkylsulfonyl.

- 10 3. A 3-(heterocyclyl)-substituted benzoylpyrazole of the formula I as claimed in claim 1 where

X is O;

- 15 R¹ is C₁-C₄-alkyl;


R⁶ is halogen, nitro, C₁-C₄-haloalkyl, C₁-C₄-alkoxy or C₁-C₄-haloalkoxy.

- 20 4. A 3-(heterocyclyl)-substituted benzoylpyrazole of the formula I as claimed in claim 1 where

X is N(C₁-C₆-alkyl).

- 25 5. A process for preparing 3-(heterocyclyl)-substituted benzoylpyrazoles of the formula I where R⁷ = hydroxyl as claimed in claim 1, which comprises acylating a pyrazole of the formula II


30




II

- 35 with an activated benzoic acid III α or a benzoic acid III β ,

40



III α




III β

- 45 where the variables X, R¹ to R⁶ and R⁸ to R¹¹ are as defined in claim 1 and L¹ is a nucleophilically replaceable leaving group and rearranging the acylation product, in the presence

50


or absence of a catalyst, to give the compounds of the formula I where $R^7 = \text{hydroxyl}$.

6. A process for preparing 3-(heterocyclyl)-substituted benzoylpyrazoles of the formula I where $R^7 = \text{OH}$ as claimed in claim 1, which comprises reacting a pyrazole of the formula II



II


in which the variables R^8 to R^{11} are as defined in claim 1, or an alkali metal salt thereof, with a 3-(heterocyclyl)benzene derivative of the formula V




V

where the variables X and R^1 to R^6 are as defined in claim 1 and L^2 is a leaving group in the presence of carbon monoxide, a catalyst and a base.

7. A process for preparing 3-(heterocyclyl)-substituted benzoylpyrazoles of the formula I where $R^7 \neq \text{hydroxyl}$ as claimed in claim 1, which comprises reacting a 3-(heterocyclyl)-substituted benzoylpyrazole I where $R^7 = \text{hydroxyl}$

I where $R^7 = \text{OH}$

with a compound of the formula VI




where

L^3 is a nucleophilically replaceable leaving group;

51

R^{7a} is C₁-C₆-alkyl, C₃-C₆-alkenyl, C₁-C₆-alkylsulfonyl, C₁-C₆-alkylcarbonyl, C₁-C₄-(alkylthio)carbonyloxy, phenylsulfonyl or phenylcarbonyl, where the phenyl radical of the two last-mentioned substituents may be partially or fully halogenated and/or may carry one to three of the following groups:
 nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy or C₁-C₄-haloalkoxy.

8. A benzoic acid ester of the formula III




where the variables X, R¹ and R³ to R⁶ are as defined in claim 1 and

R² is C₁-C₄-haloalkyl; and

L is hydroxyl or a radical that can be removed by hydrolysis.

9. A 3-(heterocyclyl)benzene derivative of the formula V



where the variables X, R¹ and R³ to R⁶ are as defined in claim 1 and

R² is C₁-C₄-haloalkyl; and

L² is a nucleophilically displaceable leaving group.

10. A composition, comprising a herbicidally effective amount of at least one 3-(heterocyclyl)-substituted benzoylpyrazole of the formula I or an agriculturally useful salt of I as

52

claimed in any of claims 1 to 4 and auxiliaries which are customarily used for formulating crop protection agents.

11. A process for preparing compositions as claimed in claim 8,
5 which comprises mixing a herbicidally effective amount of at least one 3-(heterocyclyl)-substituted benzoylpyrazole of the formula I or an agriculturally useful salt of I as claimed in any of claims 1 to 4 and auxiliaries which are customarily used for formulating crop protection agents.
- 10 12. A method for controlling undesirable vegetation, characterized in that a herbicidally effective amount of at least one 3-(heterocyclyl)-substituted benzoylpyrazole of the formula I or an agriculturally useful salt of I as claimed in
15 any of claims 1 to 4 is allowed to act on the plants, their habitat and/or on seed.
13. The use of the 3-(heterocyclyl)-substituted benzoylpyrazoles of the formula I and/or their agriculturally useful salts as
20 claimed in any of claims 1 to 4 as herbicides.

25

30

35

40

45


3-(Heterocyclyl)-substituted benzoylpyrazoles

Abstract

5

3-(Heterocyclyl)-substituted benzoylpyrazoles of the formula I

10



I

15

where:

X is O, NH or N-alkyl;

20 R¹ is alkyl;R², R³, R⁴, R⁵ are hydrogen, alkyl or haloalkyl;

25 R⁶ is halogen, nitro, haloalkyl, alkoxy, haloalkoxy, alkylthio, haloalkylthio, alkylsulfonyl or haloalkylsulfonyl;

30 R⁷ is hydroxyl, alkoxy, alkenyloxy, alkylsulfonyloxy, alkylcarbonyloxy, alkylthiocarbonyloxy, phenylsulfonyloxy or phenylcarbonyloxy, where the phenyl radical may be substituted;

R⁸, R⁹ are alkyl;35 R¹⁰ is hydrogen or alkyl;R¹¹ is hydrogen or alkyl;

and their agriculturally useful salts,

40

intermediates and processes for their preparation, and the use of these compounds or of compositions comprising them for controlling undesirable plants are described.

45

Declaration, Power of Attorney

Page 1 of 5

0050/050555

We (I), the undersigned inventor(s), hereby declare(s) that:

My residence, post office address and citizenship are as stated below next to my name,

We (I) believe that we are (I am) the original, first, and joint (sole) inventor(s) of the subject matter which is claimed and for which a patent is sought on the invention entitled

3-(Heterocyclyl)-substituted benzoylpyrazoles

the specification of which

☒ is attached hereto.

☐ was filed on _____ as

Application Serial No. _____

and amended on _____.

☒ was filed as PCT international application

Number PCT/EP99/09413

on December 2, 1999

and was amended under PCT Article 19

on _____ (if applicable).

We (I) hereby state that we (I) have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

We (I) acknowledge the duty to disclose information known to be material to the patentability of this application as defined in Section 1.56 of Title 37 Code of Federal Regulations.

We (I) hereby claim foreign priority benefits under 35 U.S.C. § 119(a)-(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or PCT International application having a filing date before that of the application on which priority is claimed. Prior Foreign Application(s)

Application No.	Country	Day/Month/Year	Priority Claimed
19936514.8	Germany	06 August 1999	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No
19855850.3	Germany	04. December 1998	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No

We (I) hereby claim the benefit under Title 35, United States Codes, § 119(e) of any United States provisional application(s) listed below.

(Application Number)

(Filing Date)

(Application Number)

(Filing Date)

We (I) hereby claim the benefit under 35 U.S.C. § 120 of any United States application(s), or § 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. § 112, I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR § 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application.

Application Serial No.

Filing Date

**Status (pending, patented,
abandoned)**

_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____

And we (I) hereby appoint **Messrs. HERBERT. B. KEIL**, Registration Number 18,967; and **RUSSEL E. WEINKAUF**, Registration Number 18,495; the address of both being Messrs. Keil & Weinkauf, 1101 Connecticut Ave., N.W., Washington, D.C. 20036 (telephone 202-659-0100), our attorneys, with full power of substitution and revocation, to prosecute this application, to make alterations and amendments therein, to sign the drawings, to receive the patent, and to transact all business in the Patent Office connected therewith.

We (I) declare that all statements made herein of our (my) own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Declaration

Page 3 of 5

0050/050555

Ulf Neidlein

NAME OF INVENTOR

Ulf Neidlein

Signature of Inventor

Date December 16, 1999

Brahmsstr.3

68165 Mannheim

Germany

Citizen of: Germany

Post Office Address: same as residence

Norbert Götz

NAME OF INVENTOR

Norbert Götz

Signature of Inventor

Date December 16, 1999

Schöfferstr.25

67547 Worms

Germany

Citizen of: Germany

Post Office Address: same as residence

Ernst Baumann

NAME OF INVENTOR

Ernst Baumann

Signature of Inventor

Date December 16, 1999

Falkenstr.6a

67373 Dudenhofen

Germany

Citizen of: Germany

Post Office Address: same as residence

Wolfgang von Deyn

NAME OF INVENTOR

Wolfgang von Deyn

Signature of Inventor

Date December 16, 1999

An der Bleiche 24

67435 Neustadt

Germany

Citizen of: Germany

Post Office Address: same as residence

5-6
Steffen Kudis
NAME OF INVENTOR

[Signature]
Signature of Inventor

Date December 16, 1999

Spelzenstr. 10
68167 Mannheim
Germany
Citizen of: Germany
Post Office Address: same as residence

Klaus Langemann
NAME OF INVENTOR

[Signature]
Signature of Inventor

Date December 16, 1999

Goldbergstr. 18
67551 Worms
Germany
Citizen of: Germany
Post Office Address: same as residence

Guido Mayer
NAME OF INVENTOR

[Signature]
Signature of Inventor

Date December 16, 1999

Gutleuthausstr. 8
67433 Neustadt
Germany
Citizen of: Germany
Post Office Address: same as residence

Ulf Mißlitz
NAME OF INVENTOR

[Signature]
Signature of Inventor

Date December 16, 1999

Mandelring 74
67433 Neustadt
Germany
Citizen of: Germany
Post Office Address: same as residence

Matthias Witschel
NAME OF INVENTOR

Matthias Witschel
Signature of Inventor

Date December 16, 1999

Wittelsbachstr. 81
67061 Ludwigshafen
Germany
Citizen of: Germany
Post Office Address: same as residence

Martina Otten
NAME OF INVENTOR

Martina Otten
Signature of Inventor

Date December 16, 1999

Gunterstr. 28
67069 Ludwigshafen
Germany
Citizen of: Germany
Post Office Address: same as residence

Karl-Otto Westphalen
NAME OF INVENTOR

Karl-Otto Westphalen
Signature of Inventor

Date December 16, 1999

Zum Pfauenturm 17
67346 Speyer
Germany
Citizen of: Germany
Post Office Address: same as residence


Helmut Walter
NAME OF INVENTOR

Helmut Walter
Signature of Inventor

Date December 16, 1999

Grünstadter Str. 82
67283 Obrigheim
Germany
Citizen of: Germany
Post Office Address: same as residence

United States Patent & Trademark Office
Office of Initial Patent Examination -- Scanning Division



Application deficiencies found during scanning:

☒ Page(s) _____ of Drawings. were not present
for scanning. (Document title)

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

☐ *Scanned copy is best available.*